





LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

---

**Eukaryotic cell division genes and their use in diagnosis and treatment of  
proliferative diseases**

---

5

In a first aspect, the present invention is related to the significant functional role of several *C. elegans* genes and of their corresponding gene products in cell division and proliferation processes that could be identified by means of RNA-mediated interference (RNAi).

10 In a second aspect, the invention relates to the identification and isolation of functional orthologues of said genes and their gene products found in other eukaryotic species, in particular man, including all biologically-active derivatives thereof.

In a third aspect, the present invention includes the use of said genes and gene products (including said orthologues) in the development or isolation of anti-proliferative agents for instance their use in appropriate screening assays and in methods for diagnosis and  
15 treatment of proliferative diseases.

In a forth aspect, the invention relates to antibodies to said gene products and their use in the development or isolation of anti-proliferative agents and in methods for diagnosis and treatment of proliferative diseases.

In a fifth aspect, the present invention is related to the use of these genes and gene products  
20 for developing structural models or other models for evaluating drug binding and efficacy as well as to any other uses which are derived from the new functions described here and which will become apparent from the disclosure of the present application for any person skilled in the art.

25 Metazoan cell division consists of an extremely complex, highly regulated set of cellular processes which must be tightly co-ordinated, perfectly timed, and closely monitored in order to ensure the correct delivery of cellular materials to daughter cells. Defects in these processes are known to cause a wide range of so-called proliferative diseases, including all forms of cancer. Since cell division represents one of the few, if not the only cellular  
30 process that is common to the aetiology of all forms of cancer, its specific inhibition has

- 2 -

long been recognised as a preferred site of therapeutic intervention. Although mitotic inhibitor drugs are recognised as one of the most promising classes of chemotherapeutic agent, screening attempts to find new drug candidates in this class have been undermined by the strong inherent tendency of such screens to identify agents that target a single  
5 protein, tubulin. Tubulin polymerises to form microtubules, the primary cytoskeletal elements needed for mitotic spindle function and chromosome segregation. Microtubule functions, however, are ubiquitously needed in almost all cell types, whether dividing or not, a fact which therefore explains many of the unwanted side effects caused by anti-tubulin drugs.

10

Perhaps the best known example of a highly successful anti-neoplastic drug that targets tubulin is provided by paclitaxel, and its marketed derivative, Taxol, from Bristol Meyers Squibb. Its applicability has indeed been seriously limited by difficulties in determining an adequate dosing regimen due to a range of problematic side effects. Taxol treatment has  
15 resulted in anaphylaxis and severe hypersensitivity reactions characterised by dyspnea and hypotension requiring treatment, angioedema, and generalised urticaria in 2-4% of patients in clinical trials. All Taxol is administered after pretreatment with corticosteroids and despite pretreatment, fatal reactions have occurred. Severe conductance abnormalities resulting in life-threatening cardiac arrhythmia occur in less than 1 percent of patients and  
20 must be treated by insertion of a pacemaker. Taxol can cause fetal harm or fetal death in pregnant women. Furthermore, administration is commonly accompanied by tachycardia, hypotension, flushing, skin reactions and shortness-of-breath (mild dyspnea).

Despite these shortcomings, Taxol has been hailed by many as the most successful new  
25 anti-cancer therapeutic of the last three decades. Clearly, there is good justification for attempting to add to the list of mitotic inhibitors used to treat cancer. However, additional drugs that target tubulin or interfere with microtubule dynamics may be expected to have similar applicability and limitations as Taxol.

- 3 -

The task of the present invention therefore is to find new potential target proteins/genes for therapeutical drugs other than tubulin that are essential for completion of mitosis. These proteins/genes may provide novel targets to screen for new anti-neoplastic or cytotoxic anti-cancer agents.

5

Unfortunately, until now, the systematic identification of such target proteins/genes using genetic screening methods has been difficult in metazoans, and has relied heavily on the use of the unicellular yeast. Several major advances in the use of certain metazoan model organisms, particularly the nematode worm *Caenorhabditis elegans*, have now begun to  
10 offer new ways of bridging this gap.

The above-mentioned task of the invention to find new potential target proteins/genes for therapeutical drugs other than tubulin involved in mitosis processes is solved by a screening assay in *C. elegans* based on 'genomic RNA mediated interference (RNAi)'  
15 combined with a highly probative microscopic assay for documenting the first rounds of embryonic cell division (Sulston *et al.*, The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev. Biol.* 100, 64-119 (1983); Gönczy *et al.*, Dissection of cell division processes in the one cell stage *Caenorhabditis elegans* embryo by mutational analysis. *J Cell Biol* 144, 927-946 (1999)). With this combination of techniques a selected  
20 gene and also a variety of selected genes can be functionally characterized with unprecedented speed and efficiency.

The nematode *C. elegans* exhibits an almost entirely translucent body throughout its development, thereby offering unparalleled microscopic access for exquisitely detailed  
25 cytological documentation, even for the earliest steps of embryogenesis. This important feature, along with its short life cycle (3-5 days), its ease of cultivation, and its low maintenance costs, has helped make *C. elegans* arguably the best studied of all metazoans. Also, sequence data are now available for over 97% of the *C. elegans* genome (*C. elegans* Sequencing Consortium. Genome sequence of the nematode *C. elegans*: a platform for  
30 investigating biology. *Science* 282, 2012-2018 (1998)). Thus, *C. elegans* has proven to be

an ideal organism for applying the new technique of RNA-mediated interference (RNAi). This technique consists in the targeted, sequence-specific inhibition of gene expression, as mediated by the introduction into an adult worm of double-stranded RNA (dsRNA) molecules corresponding to portions of the coding sequences of interest (Fire *et al.*, Potent  
5 and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391, 806-811 (1998)). For the vast majority of *C. elegans* genes tested to date, this has been shown to yield a sequence-specific inhibition of the targeted gene's expression, accompanied by clearly detectable loss of function phenotypes in the treated worm's F1 progeny (and even in some cases, in the treated worm itself).

10

A large-scale RNAi technique-based screen was performed for 2,232 (that means 96%) of the predicted open reading frames on chromosome III of *C. elegans* which is described in detail in Gönczy *et al.*, "Functional genomic analysis of cell division in *C. elegans* using RNAi of genes on chromosome III" *Nature* 408, 331-336 (2000). For the performance of  
15 this large-scale screen double-stranded RNA corresponding to the individual open reading frames was produced and micro-injected into adult *C. elegans* hermaphrodites, and the resulting embryos were analysed 24 hours later using time-lapse DIC microscopy.

Besides others, the *C. elegans* genes H38K22.2 (Genbank/EMBL ID: AL024499, provided in SEQ ID NO. 1 - 3), C02F5.1 (Genbank/EMBL ID: L14745; , provided in SEQ ID NO. 4  
20 and 5) and F10E9.8 (GenBank/EMBL ID: L10986; provided in SEQ ID NO. 6 and 7) gave rise to a phenotype detectable by the DIC-assay implying a functional role of these genes in metazoan cell division processes.

In at least one case ( for H38K22.2) it had also been possible to identify a structurally and functionally homologous gene, a so-called orthologous gene, in another species, in  
25 particular *Homo sapiens*, namely the human orthologue RP42.

For the mouse orthologue of the RP42 gene it had merely been known that the gene shows a strongly developmentally regulated expression, particularly in proliferating neuroblasts from which neocortical neurons originate (Mas *et al.*, "Cloning and expression of a novel gene, RP42, mapping to an autism susceptibility locus on 6Q16" *Genomics* 1; 65 (1), 70-  
30 74 (2000)). The functional role of RP42 in cell division and proliferation processes that

- 5 -

makes it an excellent tool for the development or identification of drugs for diagnosis and/or therapy of proliferative diseases was not known so far.

With the essential function of said genes in cell division and proliferation known, these  
5 newly identified target genes and their corresponding gene products, any homologues, orthologues and derivatives thereof represent excellent tools for use in the development and isolation of a wide range of therapeutics including anti-proliferative agents and in the development of methods for diagnosis and treatment of proliferative diseases.

10 Therefore, in a first aspect, the present invention relates to isolated nucleic acid molecules encoding a polypeptide functionally involved in cell division and proliferation or a fragment thereof and comprising a nucleic acid sequence selected from the group consisting of:

- 15 (a) the nucleic acid sequences presented in SEQ ID NO. 1 to 3, SEQ ID NO. 4 to 5, SEQ ID NO. 6 to 7, SEQ ID NO. 12 and fragments thereof and their complementary strands,
- (b) nucleic acid sequences encoding polypeptides that exhibit a sequence identity with SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 10, SEQ ID NO. 11 or SEQ ID NO. 13 of at least 25 % over 100 residues and/or which are detectable in a  
20 computer aided search using the blast sequence analysis programs with an e-value of at most  $10^{-30}$ ,
- (c) nucleic acid sequences which are capable of hybridizing with the nucleic acid sequences of (a) or (b) under conditions of medium stringency,
- 25 (d) nucleic acid sequences which are degenerate as a result of the genetic code to any of the sequences defined in (a), (b) or (c).

- 6 -

The above mentioned fragments of the isolated nucleic acid molecules may comprise a at least 15 nucleotides and preferably at least 20 nucleotides.

Additionally the above mentioned isolated nucleic acid molecules may be single or double-stranded DNA-molecules as well as single- or double-stranded RNA-molecules.

5

a):

The nucleic acid sequences of those nucleic acid molecules encoding a polypeptide functionally involved in cell division and proliferation as mentioned in a) are provided in the sequence listing

- 10 as SEQ ID NO. 1 - 3 (*C. elegans* genes H38K22.2 (Genbank/EMBL ID: AL024499)),  
as SEQ ID NO. 4 and 5 (*C. elegans* gene C02F5.1 (Genbank/EMBL ID: L14745)),  
as SEQ ID NO. 6 and 7 (*C. elegans* gene F10E9.8 (GenBank/EMBL ID: L10986)) and  
as SEQ ID NO. 12 (the human H38K22.2 orthologue, the RP42 protein (NCBI Accession No. AF292100)).
- 15 The corresponding deduced amino acid sequences of these target genes are disclosed in SEQ ID NO. 8 (for H38K22.2a), in SEQ ID NO. 9 (for H38K22.2b), in SEQ ID NO. 10 (for C02F5.1), in SEQ ID NO. 11 (for F10E9.8) and in SEQ ID NO. 13 (for RP42).

b):

- 20 Additionally, the present invention also comprises isolated nucleic acid molecules that are structurally and functionally homologous counterparts (particularly orthologues) of at least one of said target genes as disclosed in SEQ ID NO 1 to 7 or 12.

- Those homologous nucleic acid molecules may encode polypeptides that exhibit a sequence identity with SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 10, SEQ ID NO. 11 or
- 25 SEQ ID NO. 13 of at least 25 % over 100 residues, preferably of at least 30 % over 100 residues, more preferably of at least 35 % over 100 residues and most preferably at least 40 % over 100 residues.



Fig. 5 shows that the aforementioned sequence identities are significant homologies that are appropriate to identify a polypeptide as an orthologue of the target proteins as depicted in SEQ ID NO. 8 -11, and 13. Fig. 5 shows a multiple sequence alignment of the H38K22.2a family on protein level generated with a BLAST sequence analysis program. In this alignment the two *C. elegans* splice variants H38K22.2a and H38K22.2b are compared to their corresponding orthologues in *Drosophila* (CG7427), in mouse (AAF04863) and in *Homo sapiens* (AAH09478). The statistics in Fig 5 for the alignments show that the sequence identity on protein level between the *C. elegans* clone H38K22.2a and its human orthologue (AAH09478) is 36 % over 299 residues. Similarly, the sequence identities between *C. elegans* clone H38K22.2b (the other splice variant) and its human orthologue is 36 % over 238 residues. It is obvious to anyone skilled in the art that these sequence homologies are significant homologies and that therefore the human clone with the accession No. AAH09478 is unambiguously identified as the human orthologue of the *C. elegans* clones H38K22.2a and H38K22.2b.

The invention also comprises isolated nucleic acid molecules that are detectable in a computer aided search using one of the BLAST sequence analysis programs with an e-value of at most  $10^{-30}$ , preferably with an e-value of at most  $10^{-35}$ , more preferably with an e-value of at most  $10^{-40}$ .

Fig. 5 shows that the aforementioned e-values characterize significant sequence homologies that are appropriate to identify a polypeptide as an orthologue of the target proteins as depicted in SEQ ID NO. 8 -11, and 13.

The BLAST sequence analysis programs are programs used for sequence analysis that are publically available and known to anyone skilled in the art. When sequence alignments are done by a BLAST sequence analysis program, most of those programs calculate so called "e-values" to characterize the grade of homology between the compared sequences. Generally a small e-value characterizes a high sequence identity / homology, whereas larger e-values characterize lower sequence identities / homologies.

"Homology" means the degree of identity between two known sequences. As stated above, homologies, that means sequence identities, may suitably be determined by means of computer programs known in the art. The degree of homology required for the sequence variant will depend upon the intended use of the sequence. It is well within the capability

of a person skilled in the art to effect mutational, insertional and deletional mutations which are designed to improve the function of the sequence or otherwise provide a methodological advantage.

5 c):

The present invention further relates to isolated nucleic acid sequences or fragments thereof which are capable of hybridizing with the nucleic acid sequences of (a) or (b) under conditions of medium/high stringency.

10 The grade of sequence identity between a first and a second nucleic acid molecule can also be characterized by the capability of the first nucleic acid molecule to hybridize under certain conditions to a second nucleic acid molecule.

Suitable experimental conditions for determining whether a given DNA or RNA sequence "hybridizes" to a specified polynucleotide or oligonucleotide probe involve presoaking of the filter containing the DNA or RNA to examine for hybridization in 5 x SSC (sodium chloride/sodium citrate) buffer for 10 minutes, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution, 0,5 % SDS and 100 mg/ml of denaturated sonicated salmon sperm DNA (Maniatis et al., 1989), followed by hybridization in the same solution containing a concentration of 10 ng/ml of a random primed (Feinberg, A.P. and Vogelstein, B. (1983), *Anal. Biochem.* 132:6-13), <sup>32</sup>P-dCTP-labeled (specific activity > 1 x 10<sup>9</sup> cpm/μg) probe for 12 hours at approximately 45°C. The filter is then washed twice for 30 minutes in 2 x SSC, 0,5% SDS at at least 55°C (low stringency), at least 60°C (medium stringency), preferably at least 65°C (medium/high stringency), more preferably at least 70°C (high stringency) or most preferably at least 75°C (very high stringency). Molecules to which the probe hybridizes under the chosen conditions are detected using an x-ray film.

25

d):

The present invention further relates to isolated nucleic acid molecules or fragments thereof which are degenerate as a result of the genetic code to any of the sequences defined in (a), (b) or (c).

The application of automated gene synthesis provides an opportunity for generating sequence variants of the naturally occurring genes. It will be appreciated, for example, that polynucleotides coding for the same gene products can be generated by substituting synonymous codons for those represented in the naturally occurring polynucleotide sequences as identified herein. Such sequences will be referred to as "degenerate" to the naturally occurring sequences. In addition, polynucleotides coding for synthetic variants of the corresponding amino acid sequences can be generated which, for example, will result in one or more amino acids substitutions, deletions or additions. Also, nucleic acid molecules comprising one or more synthetic nucleotide derivatives (including morpholinos) which provide said nucleotide sequence with a desired feature, e.g. a reactive or detectable group, can be prepared. Synthetic derivatives with desirable properties may also be included in the corresponding polypeptides. All such derivatives and fragments of the above identified genes and gene products showing at least part of the biological activity of the naturally occurring sequences or which are still suitable to be used, for example, as probes for, e.g. identification of homologous genes or gene products, are included within the scope of the present invention.

Having herein provided the nucleotide sequences of various genes functionally involved in cell division and proliferation, it will be appreciated that automated techniques of gene synthesis and/or amplification may be used to isolate said nucleic acid molecules *in vitro*. Because of the length of some coding sequences, application of automated synthesis may require staged gene construction, in which regions of the gene up to about 300 nucleotides in length are synthesized individually and then ligated in correct succession for final assembly. Individually synthesized gene regions can be amplified prior to assembly, using polymerase chain reaction (PCR) technology. The technique of PCR amplification may also be used to directly generate all or part of the final genes/nucleic acid molecules. In this case, primers are synthesized which will be able to prime the PCR amplification of the final product, either in one piece or in several pieces that may be ligated together. For this purpose, either cDNA or genomic DNA may be used as the template for the PCR amplification. The cDNA template may be derived from commercially available or self-constructed cDNA libraries.

- 10 -

In a second aspect, the invention relates to nucleic acid probes comprising a nucleic acid sequence as previously characterized under (a) to (d) which may be a polynucleotide or an oligonucleotide comprising at least 15 nucleotides containing a detectable label.

These nucleic acid probes may be synthesized by use of DNA synthesizers according to standard procedures or, preferably for long sequences, by use of PCR technology with a selected template sequence and selected primers. In the use of the nucleotide sequences as probes, the particular probe may be labeled with any suitable label known to those skilled in the art, including radioactive and non-radioactive labels. Typical radioactive labels include  $^{32}\text{P}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ , or the like. A probe labeled with a radioactive isotope can be constructed from a DNA template by a conventional nick translation reaction using a DNase and DNA polymerase. Non-radioactive labels include, for example, ligands such as biotin or thyroxine, or various luminescent or fluorescent compounds. The probe may also be labeled at both ends with different types of labels, for example with an isotopic label at one end and a biotin label at the other end. The labeled probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs.

The invention also includes an assay kit comprising either an isolated nucleic acid molecule as defined above or a fragment thereof or a probe as defined above in a suitable container.

Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid and a certain degree of mismatch can be tolerated. Therefore, the nucleic acid molecules and probes of the present invention may include mutations (both single and multiple), deletions, insertions of the above identified sequences, and combinations thereof, as long as said sequence variants still have substantial sequence homology to the original sequence which permits the formation of stable hybrids with the target nucleotide sequence of interest.

The above identified nucleic acid molecules and probes coding for polypeptides functionally involved in cell division and proliferation or a part thereof will have a wide range of useful applications, including their use for identifying homologous, in particular orthologous, genes in the same or different species, their use in screening assays for

- 11 -

identification of interacting drugs that inhibit, stimulate or effect cell division or proliferation, their use for developing computational models, structural models or other models for evaluating drug binding and efficacy, and their diagnostic or therapeutic use for detection or treatment of diseases associated with anomalous and/or excessive cell division  
5 or proliferation, in particular neoplastic diseases, including both solid tumors and hemopoietic cancers, or coronary restenosis. Exemplary neoplastic diseases include carcinomas, such as adenocarcinomas and melanomas; mesodermal tumors, such as neuroblastomas and retinoblastomas; sarcomas and various leukemias; and lymphomas. Of particular interest are tumors of the breast, ovaries, gastrointestinal tract, liver, lung,  
10 thyroid glands, prostate gland, brain, pancreas, urinary tract, and salivary glands. Still more specific, tumors of the breast, ovaries, lung, colon, and lymphomas are contemplated.

In a third aspect, the present invention relates to the use of the above identified nucleic acid  
15 molecules and probes for diagnostic purposes. This diagnostic use of the above identified nucleic acid molecules and probes may include, but is not limited to the quantitative detection of the expression of said target genes in biological probes (preferably, but not limited to cell extracts, body fluids, etc.), particularly by quantitative hybridization to the endogenous nucleic acid molecules comprising the above-characterized nucleic acid  
20 sequences (particularly cDNA, RNA). An abnormal and/or excessive expression of said target genes involved in cell division may be diagnosed that way.

In a fourth aspect, the present invention relates to the use of the above identified nucleic acid molecules, probes or their corresponding polypeptides for therapeutical purposes.  
25

This therapeutical use of the above identified nucleic acid molecules, probes or their corresponding polypeptides may include, but is not limited to the use of said nucleic acid molecules and their corresponding polypeptides for direct or indirect inhibition of the expression of said target genes and/or for inhibition of the function of said target genes.  
30 Particularly gene therapy vectors, e.g. viruses, or naked or encapsulated DNA or RNA (e.g. an antisense nucleotide sequence) with the above-identified sequences might be suitable

- 12 -

for the introduction into the body of a subject suffering from a proliferative disease or from a disease affecting cell division for therapeutical purposes.

5 A particularly preferred therapeutical use of the above identified nucleic acid molecules or probes relates to their use in a therapeutical application of the RNAi technique, particularly in humans or in human cells.

Double-stranded RNA oligonucleotides effect silencing of the expression of gene(s) which are highly homologous to either of the RNA strands in the duplex. Recent discoveries reveal that this effect, called RNA interference (RNAi), that had been originally discovered  
10 in *C. elegans*, can also be observed in cells, particularly in human cells. Therefore the invention further comprises the use of double-stranded RNA oligonucleotides with the above identified nucleotide sequences (as stated in a) to d)), preferably with a length of at least 15 nucleotides (nt), more preferably with a length of at least 20 nt, for therapeutical silencing of the expression of genes involved in cell division or proliferation in cells of  
15 other species, particularly in human cells. This therapeutical use particularly applies to cells of an individual that suffers from a disease associated with anormalous and/or excessive cell division or proliferation, particularly a coronary restinosis or a neoplastic disease selected from the group consisting of lymphoma, lung cancer, colon cancer, ovarian cancer and breast cancer.

20

In a fifth aspect, the invention further comprises a nucleic acid construct or a recombinant vector having incorporated the nucleic acid molecules as defined in (a) to (d) or a fragment thereof.

“Nucleic acid construct” is defined herein as any nucleic acid molecule, either single- or  
25 double-stranded, in which nucleic acid sequences are combined and juxtaposed in a manner which will not occur naturally. The vector may be any vector which can be conveniently subjected to recombinant DNA procedures. The choice of the vector will usually depend on the host cell into which it is to be introduced. The vector may be an extrachromosomal entity, the replication of which is independent of chromosomal  
30 replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector in which the nucleic acid molecule as defined in (a) to (d) or a fragment thereof is operably linked to heterologous or homologous control sequences. The term "control sequences" is defined herein to include all components  
5 which are necessary or advantageous for expression of the coding nucleic acid sequence. Such control sequences include, but are not limited to, a promoter, a ribosome binding site, translation initiation and termination signals and, optionally, a repressor gene or various activator genes. Control sequences are referred to as "homologous" if they are naturally linked to the coding nucleic acid sequence of interest and referred to as "heterologous" if  
10 this is not the case. The term "operably linked" indicates that the sequences are arranged so that they function in concert for their intended purpose, i.e. expression of the desired protein.

The promoter may be any DNA sequence which shows transcriptional activity in the host  
15 cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell.

Examples of suitable promoters for directing the transcription in a bacterial host are, e.g., the phage Lambda P<sub>R</sub> or P<sub>L</sub> promoters, the *lac*, *trp* or *tac* promoters of *E. coli*, the promoter  
20 of the *Bacillus subtilis* alkaline protease gene or the *Bacillus licheniformis* alpha-amylase gene.

Examples of suitable promoters for directing the transcription in mammalian cells are, e.g., the SV40 promoter (Subramani et al., *Mol. Cell. Biol.* 1 (1981), 854-864), the MT-1  
25 (metallothionein gene) promoter (Palmiter et al., *Science* 222 (1983), 809-814) or the adenovirus 2 major late promoter.

Examples of suitable promoters for use in insect cells are, e.g., the polyhedrin promoter (Vasuvedan et al., *Febs. Lett* 311, (1992), 7-11), the *Autographa californica* polyhedrosis  
30 basic protein promoter (EP 397 485), or the baculovirus immediate early gene 1 promoter (US 5,155,037, US 5,162,222).

- 14 -

Examples of suitable promoters for use in yeast cells include promoters from yeast glycolytic genes (Hitzeman et al., *J. Biol. Chem.* 255 (1980), 1203-12080; Alber and Kawasaki, *J. Mol. Appl. Gen.* 1 (1982), 419-434) and the ADH2-4c promoter (Russell et al., *Nature* 304 (1983), 652-654).

5

The coding sequence may, if necessary, be operably linked to a suitable terminator, such as the human growth hormone terminator (Palmiter et al., *Science* 222, 809-814 (1983)), or a polyadenylation sequence. Also, to permit secretion of the expressed protein, a signal sequence may precede the coding sequence.

10

Further, the vector may comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of the plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702. Another example of such a sequence (when the host cell is a mammalian cell) is the SV40 origin of replication.

15

When the host cell is a yeast cell, suitable sequences enabling the vector to replicate are the yeast plasmid 2 $\mu$  replication genes REP 1-3 and origin of replication.

20

The vector may also comprise a selectable marker, e.g. a gene coding for a product which complements a defect in the host cell, such as the gene coding for dihydrofolate reductase (DHFR) or a gene which confers resistance to a drug, e.g. ampicillin, kanamycin, tetracyclin, chloramphenicol, neomycin or hygromycin.

25

A number of vectors suitable for expression in prokaryotic or eukaryotic cells are known in the art and several of them are commercially available. Some commercially available mammalian expression vectors which may be suitable include, but are not limited to, pMC1neo (Stratagene), pXT1 (Stratagene), pSG5 (Stratagene), pcDNA1 (Invitrogen), EBO-pSV2-neo (ATCC 37593), pBPV-1(8-2) (ATCC 37110), pSV2-dhfr (ATCC 37146).

30

In a sixth aspect, the invention comprises host cells into which the nucleic acid construct or the recombinant vector is introduced. These host cells may be prokaryotic or eukaryotic, including, but not limited to, bacteria, fungal cells, including yeast and filamentous fungi,



- 15 -

mammalian cells, including, but not limited to, cell lines of human, bovine, porcine, monkey and rodent origin, and insect cells including, but not limited to, drosophila derived cell lines.

- 5 The selection of an appropriate host cell will be dependent on a number of factors recognized by the art. These include, e.g., compatibility with the chosen vector, toxicity of the (co)products, ease of recovery of the desired protein or polypeptide, expression characteristics, biosafety and costs.

Examples of suitable prokaryotic cells are gram positive bacteria such as *Bacillus subtilis*,  
10 *Bacillus licheniformis*, *Bacillus brevis*, *Streptomyces lividans* etc. or gram negative bacteria such as *E. coli*.

The yeast host cell may be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g. *Saccharomyces cerevisiae*. Useful filamentous fungi may be selected from a species of *Aspergillus*, e.g. *Aspergillus oryzae* or *Aspergillus niger*.

- 15 Cell lines derived from mammalian species which may be suitable and which are commercially available include, but are not limited to, COS-1 (ATCC CRL 1650) COS-7 (ATCC CRL 1651), CHO-K1 (ATCC CCL 61), 3T3 (ATCCL 92), NIH/3T3 (ATCC CRL 1658), HeLa (ATCCL 2), and MRC-5 (ATCC CCL 171).

- 20 The recombinant vector may be introduced into the host cells according to any one of a number of techniques including, but not limited to, transformation, transfection, protoplast fusion, and electroporation.

- The recombinant host cells are then cultivated in a suitable nutrient medium under  
25 conditions permitting the expression of the protein of interest. The medium used to cultivate the cells may be any conventional medium suitable for growing the host cells, such as minimal or complex media containing appropriate supplements. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. in catalogues of the American Type Culture Collection).

Identification of the heterologous polypeptide expressing host cell clones may be done by several means, including, but not limited to, immunological reactivity with specific antibodies.

- 5 In a seventh aspect, the invention is related to a method for producing a polypeptide functionally involved in cell division and proliferation or a fragment thereof in a host cell comprising the steps
- (i) transferring the expression vector with an operably linked nucleic acid molecule as defined in (a) to (d) into a suitable host cell, and
  - 10 (ii) cultivating the host cells of step (i) under conditions which will permit the expression of said polypeptide or fragment thereof and
  - (iii) optionally, secretion of the expressed polypeptide into the culture medium.

In an eighth aspect, the invention comprises a polypeptide functionally involved in cell  
15 division and proliferation or a fragment thereof comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequences depicted in SEQ ID NO. 8, 9, 10, 11 and 13 and fragments thereof,
- (b) amino acid sequences which exhibit a sequence identity with the sequences of  
20 (a) of at least 25 % over 100 residues, preferably of at least 30 % over 100 residues, more preferably of at least 35 % over 100 residues and most preferably of at least 40 % over a 100 residues and/or which are detectable in a computer aided search using the BLAST sequence analysis programs with an e-value of at most  $10^{-30}$ , preferably with an e-value of at most  $10^{-35}$  and most  
25 preferably with an e-value of at most  $10^{-40}$ ,
- (c) amino acid sequences encoded by a nucleic acid molecule that is capable of hybridizing with the nucleic acid sequences of (a) or (b) or encoded by a nucleic acid molecule that is degenerate as a result of the genetic code to any of the sequences as defined in (a) or (b).

The heterologous polypeptide may also be a fusion polypeptide in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide of interest or fragment thereof. A fused polypeptide is produced by fusing a nucleic acid sequence (or a portion thereof) encoding another polypeptide to a nucleic acid sequence (or a portion thereof) of the present invention. Techniques for producing fusion polypeptides are known in the art and include ligating the coding sequences so that they are in frame and the expression of the fusion polypeptide is under control of the same promotor(s) and terminator.

10 Expression of the polypeptides of interest may also be performed using *in vitro* produced synthetic mRNA. Synthetic mRNA can be efficiently translated in various cell-free systems, including but not limited to, wheat germ extracts and reticulocyte extracts, as well as efficiently translated in cell based systems including, but not limited to, microinjection into frog oocytes, preferably *Xenopus* oocytes.

15

In a ninth aspect, the invention involves antibodies against the above identified polypeptides and against immunogenic fragments thereof. The term "antibody" as used herein includes both polyclonal and monoclonal antibodies, as well as fragments thereof, such as Fv, Fab and F(ab)<sub>2</sub> fragments that are capable of binding antigen or hapten. The present invention also contemplates "humanized" hybrid antibodies wherein amino acid sequences of a non-human donor antibody exhibiting a desired antigen-specificity are combined with sequences of a human acceptor antibody. The donor sequences will usually include at least the antigen-binding amino acid residues of the donor but may comprise other structurally and/or functionally relevant amino acid residues of the donor antibody as well. Such hybrids can be prepared by several methods well known in the art (see e.g. WO 25 89/09622; WO 94/11509; Couto, *Hybridoma* 13 (1994), 215-219; Presta, *Cancer Research* 57 (1997), 4593-4599). The antibodies of the present invention will have a wide range of useful applications, including their use for affinity purification of the corresponding immunogenic (poly)peptides, their use for the preparation of anti-idiotypic antibodies, as well as their use as specific binding agents in various assays, e.g. diagnostic or drug- 30 screening assays, or in a method for treatment of diseases associated with anomalous and/or excessive cell division or proliferation as exemplified above. Specifically, said

antibodies or suitable fragments thereof, particularly in humanized form, may be used as therapeutic agents in a method for treating cancer and other diseases associated with anomalous and/or excessive cell division or proliferation as exemplified above. Also, antibodies may be raised to the most characteristic parts of the above identified polypeptides and subsequently be used to identify structurally and/or functionally related polypeptides from other sources as well as mutations and derivatives of the above identified polypeptides.

To raise antibodies against the polypeptides of the present invention, there may be used as an immunogen either the intact polypeptide or an immunogenic fragment thereof, produced in a suitable host cell as described above or by standard peptide synthesis techniques.

Polyclonal antibodies are raised by immunizing animals, such as mice, rats, guinea pigs, rabbits, goats, sheep, horses etc., with an appropriate concentration of the polypeptide or peptide fragment of interest either with or without an immune adjuvant.

Acceptable immune adjuvants include, but are not limited to, Freund's complete adjuvant, Freund's incomplete adjuvant, alum-precipitate, water-in-oil-emulsion containing *Corynebacterium parvum* and tRNA.

In a typical immunization protocol each animal receives between about 0,1 µg and about 1000 µg of the immunogen at multiple sites either subcutaneously (SC), intraperitoneally (IP), intradermally or in any combination thereof in an initial immunization. The animals may or may not receive booster injections following the initial injection. Those animals receiving booster injections are generally given an equal amount of the immunogen in Freund's incomplete adjuvant by the same route at intervals of about three or four weeks until maximal titers are obtained. At about 7-14 days after each booster immunization or about weekly after a single immunization, the animals are bled, the serum collected, and aliquots are stored at about -20°C.

Monoclonal antibodies which are reactive with the polypeptide or peptide fragment of interest are prepared using basically the technique of Kohler and Milstein, Nature 256: 495-497 (1975). First, animals, e.g. Balb/c mice, are immunized using a protocol similar to that described above. Lymphocytes from antibody-positive animals, preferably

- 19 -

splenocytes, are obtained by removing spleens from immunized animals by standard procedures known in the art. Hybridoma cells are produced by mixing the splenocytes with an appropriate fusion partner, preferably myeloma cells, under conditions which will allow the formation of stable hybridomas. Fusion partners may include, but are not limited to:

5 mouse myelomas P3/NS1/Ag 4-1; MPC-11; S-194 and Sp 2/0. Fused hybridoma cells are selected by growth in a selection medium and are screened for antibody production. Positive hybridomas may be grown and injected into, e.g., pristane-primed Balb/c mice for ascites production. Ascites fluid is collected about 1-2 weeks after cell transfer and the monoclonal antibodies are purified by techniques known in the art. Alternatively, *in vitro*

10 production of monoclonal antibodies (mAb) is possible by cultivating the hybridomas in a suitable medium, e.g. DMEM with fetal calf serum, and recovering the mAb by techniques known in the art.

Recovered antibody can then be coupled covalently to a detectable label, such as a radiolabel, enzyme label, luminescent label, fluorescent label or the like, using linker

15 technology established for this purpose.

Antibody titers of ascites or hybridoma culture fluids are determined by various serological or immunological assays which include, but are not limited to, precipitation, passive agglutination, enzyme-linked immunosorbent antibody (ELISA) technique and radioimmunoassay techniques. Similar assays may be used to detect the presence of the

20 above identified polypeptides or fragments thereof in body fluids or tissue and cell extracts.

Assay kits for performing the various assays mentioned in the present application may comprise suitable isolated nucleic acid or amino acid sequences of the above identified genes or gene products, labelled or unlabelled, and/or specific ligands (e.g. antibodies)

25 thereto and auxiliary reagents as appropriate and known in the art. The assays may be liquid phase assays as well as solid phase assays (i.e. with one or more reagents immobilized on a support).

Unless otherwise specified, the manipulations of nucleic acids and polypeptides/-proteins can be performed using standard methods of molecular biology and immunology (see, e.g.

30 Maniatis et al. (1989), Molecular cloning: A laboratory manual, Cold Spring Harbor Lab., Cold Spring Harbor, NY; Ausubel, F.M. et al. (eds.) "Current protocols in Molecular

- 20 -

Biology". John Wiley and Sons, 1995; Tijssen, P., Practice and Theory of Enzyme Immunoassays, Elsevier Press, Amsterdam, Oxford, New York, 1985).

The invention further includes an assay kit comprising either the polypeptide as defined  
5 above or a fragment thereof or an antibody against said polypeptides as defined above or against immunogenic fragments thereof.

These recombinant polypeptides or fragments thereof as well as antibodies against those polypeptides or immunogenic fragments thereof will have a wide range of useful  
10 applications, including their use in screening assays for interacting drugs that inhibit, stimulate or effect the cell division or proliferation, their use for developing computational models, structural models or other models for evaluating drug binding and efficacy, and their use in a method for diagnosis or treatment of diseases associated with anomalous and/or excessive cell division or proliferation, in particular neoplastic diseases, including  
15 both solid tumors and hemopoietic cancers, or coronary restenosis. Exemplary neoplastic diseases include carcinomas, such as adenocarcinomas and melanomas; mesodermal tumors, such as neuroblastomas and retinoblastomas; sarcomas and various leukemias; and lymphomas. Of particular interest are tumors of the breast, ovaries, gastrointestinal tract, liver, lung, thyroid glands, prostate gland, brain, pancreas, urinary tract, and salivary  
20 glands. Still more specific, tumors of the breast, ovaries, lung, colon, and lymphomas are contemplated.

Therefore in a tenth aspect, the present invention explicitly includes the use of polypeptides as defined above or fragments thereof or of antibodies against said  
25 polypeptides or immunogenic fragments thereof in a screening assay for interacting drugs that inhibit, stimulate or effect the cell division or proliferation.

Such a screening assay for interacting drugs may particularly comprise, but is not limited to the following steps:

30

1. recombinant expression of said polypeptide or of an appropriate derivative thereof

2. isolation and optionally purification of the recombinantly expressed polypeptide or of its derivative, in particular by affinity chromatography
3. optionally labelling of the chemical compounds that are tested to interact with said polypeptide or its derivative and/or labelling of the recombinantly expressed polypeptide
4. immobilization of the recombinantly expressed polypeptide or of its derivative to a solid phase
5. binding of a potential interaction partner or a variety thereof to the immobilized polypeptide or its derivative
6. optionally one or more washing steps
7. detection and/or quantification of the interaction, in particular by monitoring the amount of label remaining associated with the solid phase over background levels.

Step 1 includes the recombinant expression of the above identified polypeptide or of its derivative from a suitable expression system, in particular from cell-free translation, bacterial expression, or baculovirus-based expression in insect cells.

Step 2 comprises the isolation and optionally the subsequent purification of said recombinantly expressed polypeptides with appropriate biochemical techniques that are familiar to a person skilled in the art.

- Alternatively, these screening assays may also include the expression of derivatives of the above identified polypeptides which comprises the expression of said polypeptides as a fusion protein or as a modified protein, in particular as a GST-fusion protein or as a protein bearing a so called "tag"-sequence. These "tags"-sequences consist of short nucleotide sequences that are ligated 'in frame' either to the N- or to the C-terminal end of the coding region of said target gene. One of the most common tags that are used to label recombinantly expressed genes is the poly-Histidine-tag which encodes a homopolypeptide consisting merely of histidines. In this context the term "polypeptide" does not merely comprise polypeptides with the nucleic acid sequences of SEQ ID No. 1 bis 7, their naturally occurring homologues, preferably orthologues, more preferably human orthologues, in particular the RP42 gene (SEQ ID No. 12), but also derivatives of these polypeptides, in particular fusion proteins or polypeptides comprising a tag-sequence.

- 22 -

These polypeptides, particularly those labelled by an appropriate tag-sequence (for instance a His-tag) or by GST, may be purified by standard affinity chromatography protocols, in particular by using chromatography resins linked to anti-His-tag-antibodies or to anti-GST-antibodies which are both commercially available. Alternatively to the use of  
5 anti-tag- or anti-GST-antibodies or other 'label-specific' antibodies the purification may also involve the use of antibodies against said polypeptides. Screening assays that involve a purification step of the recombinantly expressed target genes as described above (step 2) are preferred embodiments of this aspect of the invention.

In a third - optional - step the compounds tested for interaction may be labelled by  
10 incorporation of radioactive isotopes or by reaction with luminescent or fluorescent compounds. Alternatively or additionally also the recombinantly expressed polypeptide may be labelled.

In a forth step the recombinantly expressed polypeptide is immobilized to a solid phase, particularly (but not limited) to a chromatography resin. The coupling to the solid phase is  
15 thereby preferably established by the generation of covalent bonds.

In a fifth step a candidate chemical compound that might be a potential interaction partner of the said recombinant polypeptide or a complex variety thereof (particularly a drug library) is brought into contact with the immobilized polypeptide.

In a sixth - optional - step one or several washing steps may be performed. As a result just  
20 compounds that strongly interact with the immobilized polypeptide remain bound to the solid (immobilized) phase.

In step 7 the interaction between the polypeptide and the specific compound is detected, in particular by monitoring the amount of label remaining associated with the solid phase over background levels.

25

#### **Brief Description of the Drawings**

Fig. 1 shows DIC microscopy images taken from time-lapse recording of the first two rounds of embryonic cell division in wild type *C. elegans*.

Fig. 2 shows DIC microscopy images taken from time-lapse recording of the first two  
30 rounds of embryonic cell division in *C. elegans* F1 progeny from F0 parent treated with ds RNA "300C3" or "340G12" directed against gene H38K22.2.



Fig. 3 shows DIC microscopy images taken from time-lapse recording of the first two rounds of embryonic cell division in *C. elegans* F1 progeny from F0 parent treated with dsRNA "307C1" directed against gene C02F5.1.

Fig. 4 shows shows DIC microscopy images taken from time-lapse recording of the first two rounds of embryonic cell division in *C. elegans* F1 progeny from F0 parent treated with ds RNA "305A12" directed against gene F10E9.8.

Fig 5 shows a multiple sequence alignment of the H38K22.2a family. Herein, the amino acid sequences of the two *C. elegans* splice variants H38K22.2a and H38K22.2b are compared to the amino acid sequences of their orthologues in *Drosophila* (CG7427), in mouse (AAF04863) and in homo sapiens (AAH09478).

The "statistics" refer to values that characterize the grade of homology between the individual sequences, as the e-value, the sequence identities and the conservatively changed residues (positives).

15

#### **Description of the sequence protocol:**

- |              |  |
|--------------|--|
| SEQ ID NO. 1 | shows the unspliced DNA sequence common to both isoforms a and b of the <i>C. elegans</i> gene H38K22.2 (3104 bp). |
| SEQ ID NO. 2 | shows the spliced DNA sequence of the <i>C. elegans</i> gene H38K22.2a isoform (1011 bp).                          |
| SEQ ID NO. 3 | shows the spliced DNA sequence of the <i>C. elegans</i> gene H38K22.2b isoform (852 bp).                           |
| SEQ ID NO. 4 | shows the unspliced DNA sequence of the <i>C. elegans</i> gene C02F5.1 (3308 bp).                                  |
| SEQ ID NO. 5 | shows the spliced DNA sequence of the <i>C. elegans</i> gene C02F5.1 (3033 bp).                                    |
| SEQ ID NO. 6 | shows the unspliced DNA sequence of the <i>C. elegans</i> gene F10E9.8   |

- 24 -

- (7097 bp).
- SEQ ID NO. 7 shows the spliced DNA sequence of the *C. elegans* gene F10E9.8 (3624 bp).
- 5 SEQ ID NO. 8 shows the deduced amino acid sequence of the *C. elegans* gene H38K22.2a isoform (336 aa).
- SEQ ID NO. 9 shows the deduced amino acid sequence of the *C. elegans* gene H38K22.2b isoform (283 aa).
- SEQ ID NO. 10 shows the deduced amino acid sequence of the *C. elegans* gene C02F5.1 (1010 aa).
- 10 SEQ ID NO. 11 shows the deduced amino acid sequence of the *C. elegans* gene F10E9.8 (1207 aa).
- SEQ ID NO. 12 shows the cDNA sequence of a human orthologue of H38K22.2 (780 bp).
- SEQ ID NO. 13 shows the deduced amino acid sequence of a human orthologue of H38K22.2 (260 aa).
- 15

The following examples illustrate the present invention without, however, limiting the same thereto.

20

#### EXAMPLE 1: Generation of dsRNA molecules for RNAi experiments

First, oligonucleotide primer pair sequences were selected to amplify portions of the gene of interest's coding region using standard PCR techniques. Primer pairs were chosen to yield PCR products containing at least 500 bases of coding sequence, or a maximum of

25

- 25 -

coding bases for genes smaller than 500 bases. In order to permit the subsequent use of the PCR product as a template for *in vitro* RNA transcription reactions from both DNA strands, the T7 polymerase promoter sequence "TAATACGACTCACTATAGG" was added to the 5' end of forward primers, and the T3 polymerase promoter sequence "AATTAACCCTCACTAAAGG" was added to the 5' end of reverse primers. The synthesis of oligonucleotide primers was completed by a commercial supplier (Sigma-Genosys, UK or MWG-Biotech, Germany).

PCR reactions were performed in a volume of 50 µl, with Taq polymerase using 0.8 µM primers and approximately 0.1 µg of wild-type (N2 strain) genomic DNA template. The PCR products were EtOH precipitated, washed with 70% EtOH and resuspended in 7.0 µl TE. 1.0 µl of the PCR reaction was pipetted into each of two fresh tubes for 5 µl transcription reactions using T3 and T7 RNA polymerases. The separate T3 and T7 transcription reactions were performed according to the manufacturer's instructions (Ambion, Megascript kit), each diluted to 50 µl with RNase-free water and then combined. The mixed RNA was purified using RNeasy kits according to the manufacturer's instructions (Qiagen), and eluted into a total of 130 µl of RNase-free H<sub>2</sub>O. 50 µl of this was mixed with 10 µl 6X injection buffer (40 mM KPO<sub>4</sub> pH 7.5, 6 mM potassium citrate, pH 7.5, 4% PEG 6000). The RNA was annealed by heating at 68°C for 10 min, and at 37°C for 30 min. Concentration of the final dsRNAs were measured to be in the range of 0.1-0.3 µg/µl. The products of the PCR reaction, of the T3 and T7 transcription reactions, as well as the dsRNA species were run on 1% agarose gels to be examined for quality control purposes. Success of double stranding was assessed by scoring shift in gel mobility with respect to single stranded RNA, when run on non-denaturing gels.

25

## EXAMPLE 2: Injections of dsRNA and phenotypic assays

dsRNAs were injected bilaterally into the syncitial portion of both gonads of wild-type (N2 strain) young adult hermaphrodites, and the animals incubated at 20°C for 24 hrs. Embryos were then dissected out from the injected animals and analyzed by time-lapse

30

- 26 -

differential interference contrast videomicroscopy for potential defects in cell division processes, capturing 1 image every 5 seconds, as previously described (Gönczy *et al.*, Dissection of cell division processes in the one cell stage *Caenorhabditis elegans* embryo by mutational analysis. *J Cell Biol* 144, 927-946 (1999)). For each experiment, embryos  
5 from at least 3 different injected worms were filmed in this manner, from shortly after fertilization until the four cell stage. Embryos from 2 additional injected worms were also recorded via still images, thus yielding phenotypic documentation for at least 5 injected worms in each experiment.

In some cases, embryos exhibited acute sensitivity to osmotic changes, as evidenced by  
10 their loss of structural integrity during the dissection of the injected animals. In order to overcome this limitation, injected animals were not dissected, but rather, anaesthetized for 10 min in M9 medium containing 0.1% tricaine and 0.01% tetramisole, and mounted intact on an agarose pad to observe the F1 embryogenesis *in utero* (Kirby *et al.*, *Dev. Biol.* 142, 203-215 (1990)). The resolution achieved by viewing through the body wall does not equal  
15 that achieved by observing dissected embryos, and only limited phenotypic analysis was conducted in these cases.

Three injected animals were also transferred to a fresh plate 24 hrs after injection of dsRNA, and left at 20°C. Two days later, the plate was checked with a stereomicroscope (20-40x total magnification) for the presence of F1 larvae (L2's-L4's), as well as their  
20 developmental stage. Two days after that, the plate was inspected again for the presence of F1 adults, as well as their overall body morphology and the presence of F2 progeny.

### EXAMPLE 3: Characterization of the *C. elegans* gene H38K22.2

25

Two dsRNAs, "300C3" and "340G12", were designed and used to specifically silence the expression of the *C. elegans* gene H38K22.2 by RNAi, thereby testing its functional involvement in the first 2 rounds of embryonic cell division in this metazoan species. The dsRNAs were synthesized *in vitro* from PCR-amplified wild type genomic DNA fragments  
30 of the H38K22.2 gene. For the PCR, two sets of primer pairs were used:

- 27 -

"TCAATCAGTATGTCGACCC" with "GGAAGAAATTGGGGAAACA" as forward and reverse primers, respectively, to generate dsRNA "300C3", and "ATCGAGCGCCTCTTCAATC" with "TGGTGTCTCCATTTGCTGA" as forward and reverse primers, respectively, to generate dsRNA "340G12". The dsRNAs were purified, and injected into adult hermaphrodite worms. The phenotypic consequences of the RNAi treatment were documented 24 hours later in the F1 progeny of injected worms, using time-lapse differential interference contrast (DIC) microscopy. Embryo recordings started ~20 minutes after fertilisation, while the female pronucleus is completing its meiotic divisions, until the 4 cell stage, ~30 minutes later.

10 In the F1 progeny of control worms that were either not injected, or injected with irrelevant dsRNA, the cellular events of the first two rounds of embryonic cell division were found to exhibit very limited variability, as observed by DIC microscopy. All processes that were examined and scored for the possibility of phenotypic deviations are listed and illustrated in Figure 1. Briefly, the antero-posterior polarity of the embryo is initially determined by the position of the male pronucleus at the cortex, shortly after entry into the egg (right arrow in Fig. 1a). This is accompanied by a clear, coordinated flow of yolk granules through the central portion of the cytoplasm along the embryo's longitudinal axis towards the male pronucleus, and a concomitant series of cortical waves or ruffles progressing towards the anterior of the embryo (left side in Fig.1). Shortly thereafter, the male and female pronuclei undergo highly patterned migrations (right and left arrows respectively, in Fig. 1a,b) resulting in their meeting within the posterior half of the embryo (Fig. 1c), followed by a centration and rotation (Fig. 1d) of the pronuclear pair and associated centrosomes (arrowheads in Fig. 1b-d) to set up the future mitotic spindle along the embryo's longitudinal axis. After synchronous breakdown of the pronuclear envelopes, the clearly bipolar mitotic spindle is initially short (Fig. 1e), but then elongates while exhibiting clear lateral "rocking" movements of the posterior pole (Fig. 1f-h). These movements are accompanied by a slight posterior displacement of the posterior spindle pole, while the anterior spindle pole remains approximately stationary. This then results in an asymmetric positioning of the spindle during anaphase and telophase, thereby yielding an asymmetric placement of the cytokinetic furrow (arrowheads in Fig. 1i,j), and generating unequally-sized daughter cells: a smaller posterior P1 blastomere (right cell in Fig. 1k-o), and larger anterior AB blastomere (left cell in Fig. 1k-n). While the AB nucleus

then migrates directly to the center of the AB cell (left arrow in Fig. 1k-l), the P1 nucleus typically migrates further towards the posterior of that cell (right arrow in Fig. 1k-l), before undergoing a pronounced 90° rotation while re-migrating to the anterior P1 cortex with one of its duplicated centrosomes leading (arrowheads in Fig. 1m). This insures that the P1  
5 blastomere then divides along the embryo's longitudinal axis, perpendicular to that of the AB blastomere (Fig. 1n, arrowheads indicate centrosomes). These two divisions occur asynchronously, with P1 lagging 2-3 minutes behind AB (Fig. 1 n-p).

In the F1 embryos of worms injected with dsRNAs "300C3" or "340G12", the following highly reproducible phenotypes are observed (Fig. 2). First, although the dynamics of  
10 female pronuclear migration appear normal in all cases, its initiation is often somewhat delayed. Meeting and apposition of the two pronuclei also typically exhibits defects in that the female pronucleus gets captured by only one of the two centrosomes associated with the male pronucleus (compare Fig. 2a-c with Fig. 1a-c). Although this defect is usually corrected before pronuclear envelope breakdown is completed, subsequent positioning of  
15 the mitotic spindle within the embryo often appears defective. Weak manifestation of this phenotype appears as a lack of rocking of the posterior spindle pole during anaphase, while more severe cases show a notable drift of the entire spindle towards the posterior or lateral cortex, reaching the cortex itself and losing its longitudinal alignment completely. In the latter cases, the strongly aberrant spindle position gives rise to inappropriate specification  
20 of cleavage furrow formation, leading to anomalous cytokinesis. Even in cases where spindle position appears relatively normal, positioning of the daughter Nucleus-Centrosomes-Complexes (NCCs) typically appears abnormal as soon as anaphase ends and the cleavage furrow ingresses. This is often particularly visible in the AB blastomere, where the NCC, instead of moving directly to the centre of the cell starting at telophase,  
25 first migrates anteriorly in close proximity to the lateral cortex before eventually centering (Fig. 2a-k). This defect is usually accompanied by an apparent absence of interzonal spindle microtubules at telophase and a notable bifurcation or forking of the cytokinetic cleavage furrow (arrows in Fig. 2 g), leading to aberrantly-sized daughter blastomeres or even failure of cytokinesis by complete regression of the furrow (Fig. 2g-m). Nuclear  
30 migration and positioning of the P1 nucleus is also aberrant in most cases, resulting in a significant delay - or in some cases, a complete failure - in achieving its expected 90° rotation and association with the anterior cortex. Division of the P1 blastomere is often

significantly delayed in such embryos. Finally, defects in female meiotic divisions are also occasionally observed, as evidenced by the presence of multiple female pronuclei, indicating a failure to successfully extrude one or both polar bodies, which could come from cytokinetic defects similar to those noted above.

- 5 All observed phenotypes indicate a requirement for H38K22.2 gene function in the microtubule-dependent cellular positioning of NCCs and spindles during mitosis, and possibly meiosis. Since this function is essential to cell cycle progression and cell division throughout metazoans, this gene and any homologues and derivatives thereof represent excellent tools for use in the development of a wide range of therapeutics including anti-  
10 proliferative agents. Analysis of the H38K22.2 gene sequence reveals clear orthologues in human (NCBI Accession # AAH09478), mouse (NCBI Accession # AAF04863) and *Drosophila* (NCBI Accession # CG7427) (see Fig. 5), all of which have had no known functions ascribed to them until now. Based on their extremely high level of sequence conservation at the protein level, it can be concluded that all of these genes most likely  
15 encode proteins with equivalent functions in each of their respective species. The 336 residue protein encoded by the H38K22.2 gene isoform "a" exhibits no known structural motifs or consensus domains, according to either SMART or CDD analyses.

20 **EXAMPLE 4: Characterization of the *C. elegans* gene C02F5.1**

- A dsRNA, "307C1", was designed and used to specifically silence the expression of the *C. elegans* gene C02F5.1 by RNAi, thereby testing its functional involvement in the first 2 rounds of embryonic cell division in this metazoan species. The dsRNA was synthesized *in*  
25 *vitro* from a PCR-amplified wild type genomic DNA fragment of the C02F5.1 gene. For the PCR, oligonucleotides with sequences "ATCTGAAGATCCGTCCACT" and "ATGCACAATGGGTATTTTT" were used as forward and reverse primers, respectively, to generate dsRNA "307C1" which was purified, and injected into adult hermaphrodite worms. The phenotypic consequences of the RNAi treatment were documented 24 hours  
30 later in the F1 progeny of injected worms, using time-lapse differential interference contrast (DIC) microscopy. Embryo recordings started ~20 minutes after fertilisation,

- 30 -

while the female pronucleus is completing its meiotic divisions, until the 4 cell stage, ~30 minutes later.

In the F1 progeny of control worms that were either not injected, or injected with irrelevant dsRNA, the cellular events of the first two rounds of embryonic cell division were found to exhibit very limited variability, as observed by DIC microscopy. All processes that were examined and scored for the possibility of phenotypic deviations are listed and illustrated in Figure 1.

F1 embryos from parent worms injected with dsRNA "307C1" are consistently found to exhibit the following phenotypes (Fig. 3). First, all cellular processes that are scorable by DIC microscopy until entry into mitosis are typically indistinguishable from the wild type pattern. These include egg shape and size, yolk granule size and density, yolk granule flows and cortical ruffling, pseudo-cleavage furrow formation and positioning, pronuclear appearance (arrows in Fig. 3a) and migration (Fig. 3a,b), as well as centration and rotation of pronuclei (Fig. 3b,c) and associated pair of centrosomes (arrowheads in Fig. 3b,c). Formation and positioning of the bipolar mitotic spindle also take place normally, but the spindle is most often thinner and less rigid than in wild type, exhibiting aberrant lateral bending during its rocking and elongation at anaphase (Fig. 3f-i). After completion of cytokinesis, which appears normal, the reforming daughter nuclei are typically tear-shaped, and remain close to the newly-formed cortex for a prolonged period (Fig. 3a and k). Consistent with the tear shape, the two nuclei remain often physically connected by anomalous chromatin bridges and karyomeres are also typically seen (asterisks in Fig. 3k and l). This phenotype subsequently results in embryonic lethality in all cases.

The absence of defects in pronuclear migration and assembly of the bipolar spindle argue against a role for this gene in more general microtubule functions. The observed defects are consistent with a failure in mitotic chromosome segregation, most likely in the separation of sister chromatids, resulting in the formation of chromatin bridges, which then persist at telophase. The present data therefore indicate an essential requirement for C02F5.1 gene function in mitotic chromosome segregation. Since this function is essential to cell cycle progression and cell division throughout metazoans, this gene and any



homologues and derivatives thereof represent excellent tools for use in the development of a wide range of therapeutics including anti-proliferative agents.

Analysis of the C02F5.1 sequence reveals that the encoded 1010 residue protein contains regions predicted to form coiled coil structures, i.e. likely protein-protein interaction domains. Sequence homology analyses using the BLASTp program presently reveal no clearly orthologous sequences in other organisms. However, considering the essential and highly conserved nature of the cellular process in question, functional orthologues of this gene/protein are extremely likely to exist in all metazoans, possibly in all eukaryotes, and will be identified using for example the methodology as outlined in EXAMPLE 6.

#### **EXAMPLE 5: Characterization of the *C. elegans* gene F10E9.8**

Two dsRNAs, "305A12" and "341G5", were designed and used to specifically silence the expression of the *C. elegans* gene F10E9.8 by RNAi, thereby testing its functional involvement in the first 2 rounds of embryonic cell division in this metazoan species. The dsRNAs were synthesized *in vitro* from PCR-amplified wild type genomic DNA fragments of the F10E9.8 gene. For PCR, two sets of primer pairs were used: "TTCGTCTCGAACACGTATATCCT" with "GAAAGAAGATGAATCAGGCATTG" as forward and reverse primers, respectively, to generate dsRNA "305A12", and "CTGCAAAAATTATGACTGTGTGCG" with "AGCATTGAGATTTGGTTGTCC" as forward and reverse primers, respectively, to generate dsRNA "341G5". The dsRNA was purified, and injected into adult hermaphrodite worms. The phenotypic consequences of the RNAi treatment were documented 24 hours later in the F1 progeny of injected worms, using time-lapse differential interference contrast (DIC) microscopy. Embryo recordings started ~20 minutes after fertilisation, while the female pronucleus is completing its meiotic divisions, until the 4 cell stage, ~30 minutes later.

In the F1 progeny of control worms that were either not injected, or injected with irrelevant dsRNA, the cellular events of the first two rounds of embryonic cell division were found to

exhibit very limited variability, as observed by DIC microscopy. All processes that were examined and scored for the possibility of phenotypic deviations are listed and illustrated in Figure 1.

In the F1 embryos of worms injected with dsRNAs "305A12" or "341G5", the following highly reproducible phenotypes are observed (Fig. 4). First, all cellular processes that are scorable by DIC microscopy until the 2-cell stage are typically indistinguishable from the wild type pattern. These include egg shape and size, yolk granule size and density, yolk granule flows and cortical ruffling, pseudo-cleavage furrow formation and positioning, pronuclear appearance (arrows in Fig. 4a) and migration (Fig. 4a,b), as well as centration and rotation of pronuclei (Fig. 4b,c) and associated pair of centrosomes (arrowheads in Fig. 4b,c). The first round of division also occurs without any detectable deviations from wild type (Fig. 4d-h). It should particularly be noted that no defects are observed with respect to size, number or positioning of centrosomes or spindle poles in the single cell embryo (note arrowheads in Fig. 4b-f). In the two-cell stage embryo, however, although nuclear positioning also remains equivalent to wild type, an apparent failure in centrosome duplication is consistently observed in one of the two blastomeres and sometimes in both. A single perinuclear centrosomal region, as seen by its exclusion of yolk granules (black arrowhead in Fig. 4h-j), is typically observed instead of the two normally seen both in wild type embryos and in the unaffected blastomere (white arrowheads in Fig. 4i,j). Despite the apparent failure in centrosome duplication, microtubule-dependent processes continue normally, as illustrated by the successful anterior migration of the P1 nucleus, with its single centrosomal region leading (black arrowhead in Fig. 4h-j). Upon entering mitosis, as scored by nuclear envelope breakdown, the defective blastomere then fails to generate a bipolar spindle, forming instead a monopolar array of microtubules (dashed circle in Fig. 4k), as evidenced by the radial alignments of yolk granules in that region. Cytokinesis fails to occur in that blastomere, resulting in reformation of multiple, irregularly sized nuclei, known as karyomeres (arrows in Fig. 4m,n). In contrast, all aspects of cell division occur normally in the neighboring blastomere, resulting in normal daughter cells, each containing a single equal-sized nucleus (arrows in Fig. 4l).

The complete failure in bipolar spindle formation, accompanied by the presence of a single centrosomal region instead of two in the affected two-cell stage blastomere, clearly

indicates a requirement for F10E9.8 gene function in the complex process of mitotic spindle assembly. However, the lack of detectable defects in other microtubule-dependent processes including pronuclear migration and spindle function in the single-cell embryo effectively rules out a general microtubule-related function. In view of the maternal nature  
5 of the RNAi effect and the fact that the egg inherits its first centrosome paternally, the successful generation of a bipolar spindle in the single-cell embryo further suggests that F10E9.8 function may, in fact, be required for some aspect of centrosome duplication or separation.

Indeed, since sperm development is fully completed within the parent before initiation of  
10 the RNAi treatment, it remains unaffected by the injected dsRNA. This results in the donation of an intact "wild type" centrosome from the sperm to the egg at fertilisation. After fertilisation, this already bipartite centrosome (i.e. containing two "replication units", as evidenced by the presence of two centrioles) undergoes one round of duplication, as observed in other systems by the budding of a new centriole barrel from each existing  
15 centriole. This is followed by a physical separation of the two centriole pairs and associated pericentriolar material. This process is not dependent on the prior duplication event, and is solely needed to insure the successful formation of the bipolar spindle to be used in the first round of embryonic cell division. It therefore appears that F10E9.8 function is most likely not required for this process.

20 5. If the first duplication round fails, however, bipolar spindle formation is expected to fail during the second round of division, as seen here. Interestingly, the fact that this failure often occurs only in one of the two blastomeres suggests that in these cases only one of the original centrosome's two "replication units" actually failed in its first round of duplication at the single-cell stage. This observation is consistent with findings from other eukaryotes  
25 indicating that one of the two replication units contained within the sperm's centrosome actually comes into the egg already fully equipped for one duplication round, while the other must rely on cytoplasmic factors within the egg to permit its own duplication (Sluder, G., Hinchcliffe EH. Control of centrosome reproduction: the right number at the right time. *Biol. Cell.* 91, 413-27 (1999).

- 34 -

The present findings therefore suggest that the requirement for F10E9.8 function in mitotic spindle assembly most likely results from this gene's essential role in the process of centrosome duplication.

- 5 Since the process of spindle assembly is essential to cell cycle progression and cell division throughout metazoans, this gene and any homologues and derivatives thereof represent excellent tools for use in the development of a wide range of therapeutics including anti-proliferative agents. Analysis of the F10E9.8 sequence reveals that the encoded 1207 residue protein contains one large region predicted to form coiled coil structures, i.e. likely  
10 protein-protein interaction domains, and four predicted transmembrane domains. Sequence homology analyses using the BLASTp program presently reveal no clearly orthologous sequences in other organisms. However, considering the essential and highly conserved nature of the cellular process in question, functional orthologues of this gene/protein are extremely likely to exist in all metazoans, possibly all eukaryotes, and will be identified  
15 using for example the following methodology.

#### **EXAMPLE 6: Protocol for identifying functional orthologues in other species**

- 20 The present invention describes genes identified as having essential functions in cell division in the model organism *C. elegans*. The basis for performing research in model organisms is that the newly discovered functions for the genes in *C. elegans* will be conserved in other species including humans. Cell division is highly conserved during evolution and therefore the approach of discovering a gene function in *C. elegans* and  
25 using the information to characterise or assign functions for the human orthologue is well justified. There are two themes of conservation of genes during evolution. A gene sequence may be conserved. This means that the DNA nucleotide sequence of the gene is very similar in different species, which in turn suggests that the function of the gene is the same in the different species. As is known to any person skilled in the art, a sequence  
30 identity or homology above a particular level defines that two genes in different species code for the same gene product and gene function. Homologous genes are typically

- 35 -

identified by performing blast analysis with appropriate software, or by other approaches. For a blast search, an e-value of  $10^{-30}$  will extract the significant homologous sequences. Further phylogenetic analysis can be performed to identify which of the extracted sequences are the orthologues.

- 5 Therefore the following example for identification of orthologues can be presented. A blast search is performed using the blast sequence analysis programs and an e-value of  $10^{-3}$ . An alternative parameter can be the percentage of sequence identity. Over 100 residues, a sequence identity of 30% defines a homologous gene. After the blast search is completed, multiple sequence alignment is performed using appropriate software (for example, 10 CLUSTALW) and a neighbour joining phylogenetic tree is generated. Any person skilled in the art can identify the human orthologue from a phylogenetic tree. Essentially, the human sequence that is separated on the tree by a single speciation event or most closely related on the tree is likely to be an orthologue.

- The second theme of conservation is that the gene function can be conserved with greater 15 divergence of sequence. In the present invention this theme of conservation is not defined. However, if other genes are discovered to have functions that result in the gene product being identified as the same gene product as those claimed in the present invention then the present claims also apply to such genes.

## Claims

1. An isolated nucleic acid molecule encoding a polypeptide functionally involved in cell  
5 division and proliferation or a fragment thereof and comprising a nucleic acid  
sequence selected from the group consisting of:
  - (a) the nucleic acid sequences presented in SEQ ID NO. 1 to 3, SEQ ID NO. 4 to 5,  
SEQ ID NO. 6 to 7, SEQ ID NO. 12 and fragments thereof and their  
complementary strands,
  - 10 (b) nucleic acid sequences encoding polypeptides that exhibit a sequence identity with  
SEQ ID NO. 8, SEQ ID NO. 9, SEQ ID NO. 10, SEQ ID NO. 11 or SEQ ID NO.  
13 of at least 25 % over 100 residues and/or which are detectable in a computer  
aided search using the blast sequence analysis programs with an e-value of at most  
 $10^{-30}$ ,
  - 15 (c) nucleic acid sequences which are capable of hybridizing with the nucleic acid  
sequences of (a) or (b) under conditions of medium/high stringency,
  - (d) nucleic acid sequences which are degenerate as a result of the genetic code to any  
of the sequences defined in (a), (b) or (c).
2. A nucleic acid probe comprising a nucleic acid sequence as defined in claim 1 which  
20 may be a polynucleotide or an oligonucleotide comprising at least 15 nucleotides  
containing a detectable label.
3. A recombinant vector or nucleic acid construct having incorporated therein the  
isolated nucleic acid molecule of claim 1 or a fragment thereof.
- 25 4. The vector of claim 3 which is an expression vector.
5. A host cell which has been genetically engineered to incorporate therein the isolated  
nucleic acid molecule of claim 1 or the recombinant vector or nucleic acid construct of  
claim 3.
- 30 6. The host cell of claim 5 having incorporated therein the expression vector of claim 4.

7. An assay kit comprising the isolated nucleic acid molecule or a fragment thereof of claim 1 or the probe of claim 2 in a suitable container.
- 5 8. A method for producing a polypeptide functionally involved in cell division and proliferation or a fragment thereof in a host cell comprising the steps
  - (a) transferring the expression vector of claim 4 into a suitable host cell, and
  - (b) cultivating the host cells of step (a) under conditions which will permit the expression of said polypeptide or fragment thereof and
  - 10 (c) optionally, secretion of the expressed polypeptide into the culture medium.
9. Use of a probe as defined in claim 2 to isolate orthologues of genes comprising the nucleic acid sequences as disclosed in SEQ ID NO. 1 to 3, SEQ ID NO. 4 to 5, SEQ ID NO. 6 to 7, SEQ ID NO. 11.
- 15 10. Use of the isolated nucleic acid molecule or a fragment thereof as defined in claim 1 for producing a polypeptide functionally involved in cell division and proliferation or a fragment thereof.
- 20 11. Use of a nucleic acid molecule or a fragment thereof as defined in claim 1 or of the probe of claim 2 in a screening assay for interacting drugs that inhibit, stimulate or effect the cell division or proliferation.
- 25 12. Use of a nucleic acid molecule as defined in claim 1 or of the probe of claim 2 in a method for diagnosis or treatment of diseases associated with anormalous and/or excessive cell division or proliferation.
- 30 13. The use of claim 12 wherein the disease is a coronary restenosis or a neoplastic disease selected from the group consisting of lymphoma, lung cancer, colon cancer, ovarian cancer and breast cancer.

14. A polypeptide functionally involved in cell division and proliferation or a fragment thereof comprising an amino acid sequence selected from the group consisting of:
- 5 (a) the amino acid sequences depicted in SEQ ID NO. 8, 9, 10, 11 and 13 and fragments thereof,
- (b) amino acid sequences which exhibit a sequence identity with the sequences of (a) of at least 25 % over 100 residues and/or which are detectable in a computer aided search using the BLAST sequence analysis programs with an e-value of at most  $10^{-30}$ ,
- 10 (c) amino acid sequences encoded by any of the nucleic acid sequences (c) – (d) as defined in claim 1.
15. A fusion protein comprising the polypeptide or fragment thereof of claim 14.
- 15 16. An antibody or a fragment thereof capable of specifically binding with the polypeptide of claim 14 or with an immunogenic part thereof.
17. A humanized antibody capable of specifically binding with the polypeptide of claim 14 or with an immunogenic part thereof.
- 20 18. An assay kit comprising the polypeptide as claimed in claim 14, the fusion protein as claimed in claim 15, or the antibodies as claimed in claims 16 and/or 17 in a suitable container.
19. Use of the polypeptide of claim 14, of the fusion protein of claim 15, or of the  
25 antibodies of claims 16 or 17 in a screening assay for interacting drugs that inhibit, stimulate or effect the cell division or proliferation.
20. The use of a polypeptide or of an antibody as claimed in claim 19 wherein the screening assay for interacting drugs comprises the following steps:



1. recombinant expression of said polypeptide in a host cell
  2. isolation and optionally purification of the recombinantly expressed polypeptide of step 1
  3. optionally labelling of the drugs that are tested to interact with said polypeptide and/or labelling of the recombinantly expressed polypeptide
  4. immobilization of the recombinantly expressed polypeptide to a solid phase
  5. binding of a potential interaction partner or a variety thereof to the polypeptide
  6. optionally one or more washing steps
  7. detection and/or quantification of the interaction, in particular by monitoring the amount of label remaining associated with the solid phase over background levels.
- 
21. Use of the polypeptide of claim 14, of an amino acid sequence as defined in claim 14 or of the antibodies of claims 16 or 17 in a method for diagnosis or treatment of diseases associated with anomalous and/or excessive cell division or proliferation.
- 
22. The use of claim 20 wherein the disease is a coronary restenosis or a neoplastic disease selected from the group consisting of lymphoma, lung cancer, colon cancer, ovarian cancer and breast cancer.
- 
22. Use of the nucleic acid sequences as defined in claim 1 or the amino acid sequences as defined in claim 14 for developing computational models, structural models or other models for evaluating drug binding and efficacy.




FIG. 1




FIG. 2




FIG. 3




FIG. 4

## Multiple Sequence Alignment of the H38K22.2a family

```

CeH38K22.2a  ---MNRKLS--DQRTKLSQEVQWEQVTEAVSENFSAKANNIEYAMTLYEDNENLFAGSTP
CeH38K22.2b  ---MNRKLS--DQ-----
DmCG7427     MILONGKLSSTHEDNYKKEISLHTGEGTAFCTQONDKFLASDNYSONREYYRE--
MmAAF04863   ---MNRKLS--SQDRAVQEMIFQSSSEKTAISCSONDKLIVTDNFRONPELYIRESV
HsAAH09478   ---MNRKLS--SQDRAVQEMIFQSSSEKTAISCSONDKLIVTDNFRONPELYIRESV

```

```

CeH38K22.2a  QPSRPSNTEREFNQYVDEKQVGEKRMCPHGSENRLLTDEGYEATDRRLVAAVHETAT
CeH38K22.2b  ---KTKLRRFNFQYVDEKQVGEKRMCPHGSENRLLTDEGYEATDRRLVAAVHETAT
DmCG7427     ---KTKLRRFNFQYVDEKQVGEKRMCPHGSENRLLTDEGYEATDRRLVAAVHETAT
MmAAF04863   KGSERKKEQYTRKKEBOI---ENKSIDEQQFCDDALDPASISMLLANKGERAT
HsAAH09478   KGSERKKEQYTRKKEBOI---ENKSIDEQQFCDDALDPASISMLLANKGERAT

```

```

CeH38K22.2a  QGKESLDWVKETATQAPTQVNTQRDSINSGRSEKATHEHLVLAANYKSAACRN
CeH38K22.2b  QGKESLDWVKETATQAPTQVNTQRDSINSGRSEKATHEHLVLAANYKSAACRN
DmCG7427     QGKESLDWVKETATQAPTQVNTQRDSINSGRSEKATHEHLVLAANYKSAACRN
MmAAF04863   QGKESLDWVKETATQAPTQVNTQRDSINSGRSEKATHEHLVLAANYKSAACRN
HsAAH09478   QGKESLDWVKETATQAPTQVNTQRDSINSGRSEKATHEHLVLAANYKSAACRN

```

```

CeH38K22.2a  TDLHTACCQDVFQGSTIMTQIDELWAQENAAASRLAQNVGASNAKQFKSVWISRET
CeH38K22.2b  TDLHTACCQDVFQGSTIMTQIDELWAQENAAASRLAQNVGASNAKQFKSVWISRET
DmCG7427     TDLHTACCQDVFQGSTIMTQIDELWAQENAAASRLAQNVGASNAKQFKSVWISRET
MmAAF04863   TDLHTACCQDVFQGSTIMTQIDELWAQENAAASRLAQNVGASNAKQFKSVWISRET
HsAAH09478   TDLHTACCQDVFQGSTIMTQIDELWAQENAAASRLAQNVGASNAKQFKSVWISRET

```

```

CeH38K22.2a  WNFWDHILLSKPDLDYUDEGAWVLTQVNYCRENLNYPKPGNASNDQOMETPKIAQ
CeH38K22.2b  WNFWDHILLSKPDLDYUDEGAWVLTQVNYCRENLNYPKPGNASNDQOMETPKIAQ
DmCG7427     WNFWDHILLSKPDLDYUDEGAWVLTQVNYCRENLNYPKPGNASNDQOMETPKIAQ
MmAAF04863   WNFWDHILLSKPDLDYUDEGAWVLTQVNYCRENLNYPKPGNASNDQOMETPKIAQ
HsAAH09478   WNFWDHILLSKPDLDYUDEGAWVLTQVNYCRENLNYPKPGNASNDQOMETPKIAQ

```

```

CeH38K22.2a  KKPGIFYFNSNLQIEFKLFQYPMKLTIFKTIHTAGTNR
CeH38K22.2b  KKPGIFYFNSNLQIEFKLFQYPMKLTIFKTIHTAGTNR
DmCG7427     SQRNISSAYQTSHTNMNYG-----
MmAAF04863   -----
HsAAH09478   -----

```

Statistics	HsAAH09478	MmAAF04863	DmCG7427
CeH38K22.2a	E-value: 1e-49 Identities: 101/275 (36%) Positives: 158/275 (56%)	E-value: 7e-49 Identities: 100/275 (36%) Positives: 157/275 (56%)	E-value: 6e-44 Identities: 104/299 (36%) Positives: 154/299 (56%)
CeH38K22.2b	E-value: 1e-35 Identities: 118/275 (43%) Positives: 180/275 (65%)	E-value: 7e-35 Identities: 117/275 (43%) Positives: 177/275 (65%)	E-value: 1e-36 Identities: 126/299 (42%) Positives: 179/299 (60%)

FIG. 5

CE61773US.ST25  
SEQUENCE LISTING

<110> Cenix BioScience GmbH

<120> Eukaryotic cell division genes and their use in diagnosis and treatment of proliferative diseases

<130> CE61773US

<150> US 60/246,750

<151> 2000-11-09

<160> 26

<170> PatentIn version 3.1

<210> 1

<211> 3104

<212> DNA

<213> C. elegans

<400> 1

```

atgaatcgac tgaagtcgac tcaaaaaaca aaggtttgta aacggaaaca agacgatgaa   60
gtggagatga gtgatatgga aactgacac aaaaagtgta gaaaacaaga aaacagtaaa  120
tttgtgctgt tgaaaattcc attcgatcac cattcccggt tttctctttt tcagcattta  180
tctcgagcaa gttcgagttc tctagctcaa agcactgttc tttctgacat ttttcccaag  240
aactacgata atatcgtgag ttgtagcggg aatttcgaaa aaaaaactaa ttttgccaca  300
tcttgctgct tcgtttgtta tttcttgact agacaaattc tagctcatct agaaagctga  360
cttttctcaa aatcgttgcg agaccctaaag cagaaaaatg tatctttttt aaatctacgt  420
ggaaacgcgc tccaatatta aatttcgagg ttttcccgcc aaatacctaa cgagaccctaa  480
ctttggcgag cagagcggtt tgcccgcgat tttcctgcgt ctcttcaaac aatctaata  540
ctgctgctgg tttatgaaat atcaattttc ctcatTTTTT aaagctgagc aatgttttcg  600
ctcaatccta aaatttttag tagttctaata tgtgatcaac ggtttcccat ttccgatcga  660
agtcactttt taaattctca cttttattga tttttttcgt tttgaaattc ctgatttctt  720
ccttttttagt gataagacat cagttgctga ctgtagagaa agtgtgagaa actgttagtg  780
agagagagaa aacagtttga gaaaatgaaa aatgttttaa ataatagatat cataattatt  840
atttgatacc atttccagct ccggcagttc gtccagtgga ctcagggtcac ggaagctgtg  900

```

## CE61773US.ST25

tctctcaact tcctggcaaa agctaattgg aatatcgaat acgcatgac tctgtatttc 960  
gacaatccta atctttttgc tggatcgaca ccacagccga gcgttgatag gtccaatgta 1020  
cggcaattgc tgactttggc aactctacag aatgataatg ttctcacaat atttttaatt 1080  
aaaatttagt tatatttaga ctatagaaaa aatatttgat ttatctgaaa atacatttta 1140  
tttcagttgg aataattgga aaagtgtctt caaataattg tttttgagcg cttttttaat 1200  
tgttccaact gaaatcaaag ccattttcag ataaagcaaa tttttttaaa gtatatcact 1260  
aagttttaat tctaaaaaag tattgggaga acatgtcaca ccgactcatt ttgttgaatt 1320  
gccgacaatt gcagaattta aatttaatta tgtaaataaa agtaattttt gtagatcgag 1380  
cgcctcttca atcagtatgt cgacccaaag gataaagttg gagaaaaacg aatgggaccc 1440  
cacggaatca atcgtttgct cactgatctt ggctatgaag ctactgatcg ccgggttctt 1500  
gtgctcgctt ggaagtgttac tgcacagaca caatgtgaat tctcgttgga tgaatgggtg 1560  
aaaggaatga cagctcttca agcggatact gttcaaaatt tgagacaacg aatcgattcg 1620  
attaattcag gactggaatc ggataaggca aaagtacgga aaaaattaaa taactggaat 1680  
tatcttccaa acttatttga aagtgggaga gcgaatttgc actttttaag aacaaattca 1740  
cgcaaaacac tgtaaattga agttaattga aaaattttga tgtaaatac agagaaaaat 1800  
tacacacttt tcctcgagga gtacacgggc tgcgtaaatc aacacatagc tttattgttg 1860  
gttcacacca cggcagtatg ataatcaaaa aaaaaattta attgaaaaat tgaaattaag 1920  
atggaggaaa atgttatttc gatctggaaa taatatttat ttttgatgaa attaataaat 1980  
ataattttca gaccgaagga aaattttaat acgtttctat aataattttc gattcaaaaa 2040  
tttgaattat cacaattttt aaaaacaaaa aggttctacg atcgtctcat atctaataatc 2100  
ttatcagtta cagttccacg agctctacct attgccttc aactatgcca aatccgccgc 2160  
ttgccgcaat ctggatcttg aaactgcat ctgttgctgg gatgttcttt tcggacaacg 2220  
atcaacaatt atgactcaat ggatcgattt tctatgggca caggagaacg cggcggcgtc 2280  
tcgcctcgct cagaacgtgg gcgcttccaa tgcgaagcaa ttcaaactcg tgtggatctc 2340  
tcgtgacacg tggaatctct tctgggactt tattcttctg agtaagccag atttgctgga 2400  
ttacgatgat gaaggagcat ggccagtgtt tattgatcaa ttcgttgatt attgccgtga 2460



## CE61773US.ST25

aaatctcaat tatccaaagc caggaaatgc gtcaaatgat cagcaaattg agacaccaag 2520  
 ttattattag gacaaacaat tctaaaatcc taaagggttcg tgtttcccca atttcttcct 2580  
 attttcagag ttataaaata ttgcctggac gcgaaatfff gcttcaaaac tacggtacca 2640  
 ggtctcggca cgacaaatat tggttaaatg cgaaaatgca cgcgcttca atgggtactg 2700  
 tagtttcaca cttttcaaaa cgtaattttt tctatgacaá cagataagct ttaaaaaatc 2760  
 ttgtgaaaaa cttcaaaaaa tcaaaagtff gaaggcgac atattttaac aaaaaatgff 2820  
 tcgtgccgag accggctacc gtatttttta tgcgaaatff cgcgtttgtg taatattttt 2880  
 atattatacc gagaaaactc gacactttaa aggtgtggta gcgaattggg attttatttc 2940  
 gaaaaatatc ctaaatattc ccaaattcag aatagcgca aaagaaaccg ggaatttttt 3000  
 attttaattc taatttaca ctaatagaat tcaaattgff tcagtatccc atgctcaaga 3060  
 ctatcttcaa aataacaatt cacaccgccg gaacaaatcg ataa 3104

<210> 2  
 <211> 1011  
 <212> DNA  
 <213> C. elegans

<400> 2  
 atgaatcgac tgaagtccga tcaaaaaaca aagctccggc agttcgtcca gtggactcag 60  
 gtcacggaag ctgtgtctct caacttcctg gcaaaagcta attggaatat cgaatacgcg 120  
 atgactctgt atttcgacaa tcctaattctt ttgtctggat cgacaccaca gccgagcgff 180  
 gataggcca atatcgagcg cctcttcaat cagtatgtcg acccaaagga taaagttgga 240  
 gaaaaacgaa tgggaccca cggaatcaat cgtttgtcga ctgatcttg ctatgaagct 300  
 actgatcgcc gggttcttgt gctcgcttg aagtttactg cacagacaca atgtgaattc 360  
 tcgttgatg aatgggtgaa aggaatgaca gctcttcaag cggatactgt tcaaaatttg 420  
 agacaacgaa tcgattcgat taattcagga ctggaatcg ataaggcaaa attccacgag 480  
 ctctacctat ttgccttcaa ctatgccaaa tccgccgctt gccgcaatct ggatcttgaa 540  
 actgccatct gttgctggga tgttcttttc ggacaacgat caacaattat gactcaatgg 600  
 atcgattttc tatgggcaca ggagaacgag cggcgctctc gcctcgctca gaacgtgggc 660  
 gcttccaatg cgaagcaatt caaatcgggtg tggatctctc gtgacacgtg gaatctcttc 720

## CE61773US.ST25

```

tgggacttta ttcttctgag taagccagat ttgtcggatt acgatgatga aggagcatgg 780
ccagtgccta ttgatcaatt cgttgattat tgccgtgaaa atctcaatta tccaaagcca 840
ggaaatgcgt caaatgatca gcaaatggag acaccaaaaa tagcgcaaaa gaaacccgga 900
atTTTTtatt ttaattctaa ttacaacta atagaattca aattgtttca gtatcccatg 960
ctcaagacta tcttcaaaat aacaattcac accgccggaa caaatcgata a 1011

```

```

<210> 3
<211> 852
<212> DNA
<213> C. elegans

```

```

<400> 3
atgaatcgac tgaagtccga tcaaaaaaca aagatcgagc gcctcttcaa tcagtatgtc 60
gacccaaagg ataaagttgg agaaaaacga atgggacccc acggaatcaa tcgtttgctc 120
actgatcttg gctatgaagc tactgatcgc cgggttcttg tgctcgctg gaagtttact 180
gcacagacac aatgtgaatt ctcgttggat gaatgggtga aaggaatgac agctcttcaa 240
gcgatactg ttcaaaattt gagacaacga atcgattcga ttaattcagg actggaatcg 300
gataaggcaa aattccacga gctctaccta tttgccttca actatgccaa atccgccgct 360
tgccgcaatc tggatcttga aactgccatc tgttgctggg atgttctttt cggacaacga 420
tcaacaatta tgactcaatg gatcgatttt ctatgggcac aggagaacgc ggcggcgtct 480
cgctcgctc agaacgtggg cgcttccaat gcgaagcaat tcaaatcggg gtggatctct 540
cgtgacacgt ggaatctctt ctgggacttt attcttctga gtaagccaga tttgtcggat 600
tacgatgatg aaggagcatg gccagtgctt attgatcaat tcgttgatta ttgccgtgaa 660
aatctcaatt atccaaagcc aggaaatgcg tcaaatgatc agcaaagga gacaccaaaa 720
atagcgcaaa agaaacccgg aatTTTTtatt ttaattcta atttacaact aatagaattc 780
aaattgtttc agtatcccat gctcaagact atcttcaaaa taacaattca caccgccgga 840
acaaatcgat aa 852

```

```

<210> 4
<211> 3308
<212> DNA

```

CE61773US.ST25

&lt;213&gt; C. elegans

&lt;400&gt; 4

```
atgtcgatgg agcctcgtaa gaagcggaac tcgattctca aggtgcggca agccgctcgaa 60
accatcgagg aaaccgtcat gaacagtggg cctagttcca caacaactaa tcgacgagtc 120
agctttcata acgtgaagca tgtcaagtca gttagagtca gtgaataatt tatcaataaa 180
ataattatit caggcagtat gacagggacc atggtaaaat tcttgacgcc acaccagtta 240
aggagaagat tactgacact attggatcag atggtatitit gacgtgagtt ccatccttta 300
acgtgaaata atgaatacgt aaaaatctit ttaagaccac gtggcggaag catggatatt 360
tccgaatctc cggcctgcac gtcctcattt caagtgttcg gcggtggtaa tctcgataaa 420
actatggata tgtctctcga aacaactatc aacgagaaca acgaaacggc gagattgtit 480
gaaaccacaa gagatccaac actattatac gaaaagatcg tcgaaaccac aacaaaagtt 540
accgagcgaa ttgttagtat gccactggat gataccttag caatgttcaa tacaacgaat 600
caagaagata aggatatgtc agttgatcgt tcagttctit tcacgattcc caaagttccg 660
aagcataacg ctacaatgaa tagaactata ccgatggacc tcgatgaatc aaaagcagcg 720
ggcgggccagt gcgatgaaac ggtatgttga attaatagaa ggaaccaaait tatcttaait 780
ttacagatga atgtgttcaa ttccacaaac ttggaagccg ctgaaatgga tacgagtaaa 840
ttagatgaaa ataataccat gaatgctatc cggattccga ttaaitcaaa cgtcatgcct 900
gtagacatgg acatcactga acatcacact ttaaitgaag aaaagaaaaa tgatacattc 960
gggccaagtc aactgatgga catttcggcg ccacaagttc aagttaatga tactttggcc 1020
atititcaaca gtccgagaga catctgtaait aagggtitgg gtgttctca gaatctaata 1080
aatatgcctc cgaacgtcgt acctgtggac atggacatca ctgatcaggc cgtattaaac 1140
gcggagaaga aaaaatgatca attcgagaca agtcagctta tggacatttc tattccgaaa 1200
gttctagtaa atgacactat ggcgatgttc aacagcccga aacacgtcag taagagcagc 1260
atggatctcg agaaaacgat tgaagccgct gacaaatcaa cgaaataccc gagtatcgca 1320
gatgaggtgg aagatttaga catggatatg gatatcactg aacaacaacc atgtgaggct 1380
ggtaatcagc agaacgacgg cttgcaactt caaaaggagg atttaatgga catttcgggtg 1440
attcgagatt cacctgcagt aaacgacacc atggctgtgt tccagagtcc tgccagagta 1500
```

## CE61773US.ST25

aagatcggag cggttaagttt taagcacact ttccaataaa aatgtatttc tttcagaaca 1560  
actcgatcat tgattcgcag aaatctatcg tgttcgggtga cgaaatgagc attgacgaga 1620  
cacaaaatga tggaaccttg acgttgccaa agtcgaatgt agaagtgact acaactaatg 1680  
atgtctacac gtctctcgag cggcaagagg aaaatgcttc agaaaacgta tccatgataa 1740  
acgaatcttc tgttcattcg gaaatcgaca aaaagtcgtt tatgctcatc gaagaagaaa 1800  
gggcttttat gcaactcctcc atgattgatg tagcacaaaa gttggaagac gatggttcgt 1860  
cgaagacgcc agtcacacctt gcttcacagt cagcttctct tgccactaaa gaaccatcag 1920  
cccttcacaa ctcgagtgc actctcaaca attcgatgga attggacaac aatactcttc 1980  
ttaaaactat gcaaattaca acgtgtgaag acattagcat ggtccatgag tctattgctg 2040  
ttgaactgaa cagtaacaaa gagcaggagc aattcggaga tgagactttg cagaaaaatg 2100  
gtaaatttcg tttattcaat aactctatta aaagtatgtt ttagatacct cgaatactgg 2160  
cgcgaaatttc acattccaag gccataatga aacatcgcaa atcatgaaca atgtcgactc 2220  
ggaagcagtg aacacgtcca agatttcaac atattcggct ttcaatttga gcatcaacca 2280  
gtctatctct aaacgacgtc gatctcttct gaattctgct cgtgaatctc ctgcgtcgtg 2340  
tgcgttggag aattctataa tgtcgatgaa tgggcaaaca atggaagctc tgacagaata 2400  
tcgacagaat aaaactatgc agacgagtc agattcgatg ccgagtatga gtttgaacga 2460  
ttcgggaaga gatattctcg cgatggtaag aatatctctt tgagtattga atcgaaaatg 2520  
tctttcagaa tacatcagtc cgctctctc atctgaattc ttcaaaaact gctgccccag 2580  
gaacaccatc attgatgtca caaatgtac aacttcacc tccatctcct caattcgaaa 2640  
tgccagactt cgatccagct gtggtcaacg ttgtatattt aacatctgaa gatccgtcca 2700  
ctgaacaaca tccagaagct ctcaaatttc agcgtattgt tgaaaacgag aaaatgaaag 2760  
tacaacacga gattgattct ctgaattcaa ccaatcaact ttctgctgag aaaattgata 2820  
tgttgaagac taaggagctc ttgaagttta gtcatgatga gcgagaagcg attatgattg 2880  
caagaaaaga cgcggaaatc aagtttttgg agcttcgtct gaaatttgca ctcgagaaaa 2940  
aaattgaaag tgaccaggaa attgctgaac tagaacaagg aaattcgaaa atggctgagc 3000  
agctaagagg tctcgataag atggctgtcg ttcaaaaaga actagaaaag ctgagaagtc 3060

## CE61773US.ST25

ttcctccatc acgcgaagag agcgggaaaa tccgaaagga gtggatggag atgaagcaat 3120  
 gggaattcga ccagaaaatg aaagcactcc gaaatgtacg ctcaaacatg attgcacttc 3180  
 gttcagagaa aaatgctctc gaaatgaaag tcgcggaaga acacgagaag tttgcccaga 3240  
 ggaacgattt gaagaaaagt cgaatgctgg tgttctctaa ggctgttaag aaaattgtga 3300  
 acttctag 3308

<210> 5  
 <211> 3033  
 <212> DNA  
 <213> C. elegans

<400> 5  
 atgtcgatgg agcctcgtaa gaagcggaaac tcgattctca aggtgcggca agccgctgaa 60  
 accatcgagg aaaccgtcat gaacagtggg cctagttcca caacaactaa tcgacgagtc 120  
 agctttcata acgtgaagca tgtcaagcag tatgacagg accatggtaa aattcttgac 180  
 gccacaccag ttaaggagaa gattactgac actattggat cagatggat tttgacacca 240  
 cgtggcggaa acatggatat ttccgaatct ccggcctgca cgtcctcatt tcaagtgttc 300  
 ggcggtggta atctcgataa aactatggat atgtctctcg aaacaactat caacgagaac 360  
 aacgaaacgg cgagattgtt tgaaaccaca agagatccaa cactattata cgaaaagatc 420  
 gtcgaaacca caacaaaagt taccgagcga attgttagta tgccactgga tgatacctta 480  
 gcaatgttca atacaacgaa tcaagaagat aaggatatgt cagttgatcg ttcagttctt 540  
 ttcacgattc ccaaagttcc gaagcataac gctacaatga atagaactat accgatggac 600  
 ctcgatgaat caaaagcagc gggcggccag tgcgatgaaa cgatgaatgt gttcaatttc 660  
 acaaacttgg aagccgctga aatggatacg agtaaattag atgaaaataa taccatgaat 720  
 gctatccgga ttccgattaa ttcaaacgtc atgcctgtag acatggacat cactgaacat 780  
 cacactttta ttgaagaaaa gaaaaatgat acattcgggc caagtcaact gatggacatt 840  
 tcggcgccac aagttcaagt taatgatact ttggccattt tcaacagtcc gagagacatc 900  
 tgtaataagg gtttgggtgt tcctcagaat ctaataaata tcgcctcgaa cgtcgtacct 960  
 gtggacatgg acatcactga tcaggccgta ttaaacgcgg agaagaaaaa tgatcaattc 1020

## CE61773US.ST25

gagacaagtc agcttatgga catttctatt ccgaaagttc tagtaaatga cactatggcg 1080  
atgttcaaca gcccgaaaca cgtcagtaag agcagcatgg atctcgagaa aacgattgaa 1140  
gccgctgaca aatcaacgaa atacccgagt atcgcagatg aggtggaaga tttagacatg 1200  
gatatggata tcaactgaaca acaaccatgt gaggttggtta atcagcagaa cgacggcttg 1260  
caacttcaaa aggaggattt aatggacatt tcggtgattc gagattcacc tgcagtaaac 1320  
gacaccatgg ctgtgttcca gagtcctgcc agagtaaaga tcggagcgaa caactcgatc 1380  
attgattcgc agaaatctat cgtgttcggt gacgaaatga gcattgacga gacacaaaat 1440  
gatggaacct tgacgttgcc aaagtcgaat gtagaagtga ctacaactaa tgatgtctac 1500  
acgtctctcg agcggcaaga ggaaaatgct tcagaaaacg tatccatgat aaacgaatct 1560  
tctgttcatt cggaatcga caaaaagtcg tttatgctca tcgaagaaga aagggtttt 1620  
atgcactcct ccatgattga tgtagcaciaa aagttggaag acgatgggtc gtcgaagacg 1680  
ccagtcatcc ttgcttcaca gtcagcttct cttgccacta aagaaccatc agcccttcac 1740  
aactcgagtg caactctcaa caattcgatg gaattggaca acaatactct tcttaaaact 1800  
atgcaaatta caacgtgtga agacattagc atggtccatg agtctattgc tgttgaactg 1860  
aacagtaaca aagagcagga gcaattcgga gatgagactt tgcagaaaaa tgatacctcg 1920  
aatactggcg cgaatttcac attccaaggc cataatgaaa catcgcaaat catgaacaat 1980  
gtcgactcgg aagcagtga cactccaag atttcaacat attcggcttt caatttgagc 2040  
atcaaccagt ctatctctaa acgacgtcga tctcttctga attctgctcg tgaatctcct 2100  
cgtcgtgttg cgttgagaa ttctataatg tcgatgaatg ggcaaacaat ggaagctctg 2160  
acagaatate gacagaataa aactatgcag acgagtcaag attcgatgcc gagtatgagt 2220  
ttgaacgatt cggaagaga tattctcgcg atgaatacat cagtccgctc tcctcatctg 2280  
aattcttcaa aaactgctgc cccaggaaca ccatcattga tgtcacaaaa tgtacaactt 2340  
ccacctccat ctctcaatt cgaaatgcc gacttcgatc cagctgtggt caacgttgta 2400  
tatttaacat ctgaagatcc gtccactgaa caacatccag aagctctcaa atttcagcgt 2460  
attgttgaaa acgagaaaat gaaagtacaa cacgagattg attctctgaa ttcaaccaat 2520  
caactttctg ctgagaaaat tgatatgttg aagactaagg agctcttgaa gtttagtcat 2580

## CE61773US.ST25

gatgagcgag aagcgattat gattgcaaga aaagacgcgg aaatcaagtt tttggagctt 2640  
cgtctgaaat ttgcactcga gaaaaaaatt gaaagtgacc aggaaattgc tgaactagaa 2700  
caaggaaatt cgaaaatggc tgagcagcta agaggtctcg ataagatggc tgtcgttcaa 2760  
aaagaactag aaaagctgag aagtcttcct ccatcacgcg aagagagcgg gaaaatccga 2820  
aaggagtgga tggagatgaa gcaatgggaa ttcgaccaga aaatgaaagc actccgaaat 2880  
gtacgctcaa acatgattgc acttcgttca gagaaaaatg ctctcgaaat gaaagtcgcg 2940  
gaagaacacg agaagtttgc ccagaggaac gatttgaaga aaagtcgaat gctgggtgttc 3000  
tctaaggctg ttaagaaaat tgtgaacttc tag 3033

<210> 6  
<211> 7097  
<212> DNA  
<213> C. elegans

<400> 6  
accgcatctc ttccaatgga tcaaccatca ttgtcatctt cgccggaaaa tcgtctaaat 60  
cccgcacctt ccgttgctga agagcatggc cacagtggac agcacgctga agaagaagaa 120  
gacaatgaca cggatgaagt atctgcaatg ccttcttttg tgctgatga acettcgact 180  
cttgtaatt cagatcatga attgtctgat gatgctttaa agtataaaaa tgcagctgcc 240  
gaattcaaag cttttgagag aagaatggat tcggtaagaa cagccaaatc agaatgataa 300  
ttgaaatttt acatagaata gatttacgta tcaaaaatca aaacctacga atactctcta 360  
attcaaaatt taattaatta aaattaaaga tgagatcagc ttcaacaatc acaacatcac 420  
tggcaacgcc atcatcttgt gcaccatcaa actcctctga gcctcctact cggctctacac 480  
caattatgaa cgatttaggc gttggcccaa ataatcacia ttggccgtct tcaatgcaag 540  
aattatcagg aatttctctg gaaacaccac aggctcgacc gcttggcagc aatagaatta 600  
atcagcttgg taggttaata acaaaaaaaaa catgattgat tagattttta gttcgaagtg 660  
aggctcaaac gggaataagc cttttacaac accatgaaag acctactgtg accgccccat 720  
tgagacgaaa tgatatgatg aactcatcac gacagaatcc acagaatgga aatgttcaag 780  
atgaaaatcg acccgagcac gtttatgatc aaccaatata tgttcctgga tcatcactgg 840  
accgacagaa acttgaaatt gaaattcgac gtcatcgtaa cttgaacata caactgagag 900

## CE61773US.ST25

aactctgccc gaaccgtgga cagtggaata tcaaatgaag acgagacccg tccaccaacc 5640  
aatagctaac ggtcttcttt tcgaatattc caatggagat cttcgatggg ttaatcggca 5700  
gaacgctggt aatgtaagtt ttaattggaa ttttgtcaat taaagtgacc aatttacaga 5760  
tctacatatc cgcagttgat aaaacagtca gaattgatct cccacatac aatatttcaa 5820  
ttattcatac atttcaaagg caagttgaag tacttcgtcc tggaaataac ataacattga 5880  
taagtattaa acgacgagaa gttcgaactg atttgattta tcaaaacgga atgtataaaa 5940  
ctgaaatgta agattatttt ctttttttaa gttcatcgga aatttcgtat ttcagcttca 6000  
atagggacgg aagatatggt acgaaggatt ttagcaatca agaagtttcg agaaagtgag 6060  
tctctattca ttttccaatt aattattcag aaaaaccatt aaaatctcaa aactattacc 6120  
cagatgttta atttaattta atttaattta ttacgataga agatatcgtt aggtagaaaa 6180  
aaaaacacac acacattaat agatacaaac catcacaagt ggttacataa ataaattaca 6240  
taaataaaac gaaacaaaaa taaaaaaaga gatgtgacat tttgcggcaa aaaatgtctc 6300  
ggcacgataa aatttagtta aatgggaaaa ggcgtgcgcc tttaaatatt actgtagttt 6360  
aaaaatcgcg ttactgtcga attgttgttt gcccttttt tttttgataa aaacatgttt 6420  
attagtttag aaaaaagata aataaaccaa actacaacag tctttatagg cgcacgtatt 6480  
ttcacattta aaaatctgtc ctttaacgaa aaaattgtaa aatttggcgc cttcaaagag 6540  
tactgtaatt tcaaactcaa tttgaaacag aattttcatc gattttcctt agttagtttt 6600  
tcgatgaatt ttaatttatt cattaaaaaa actcaaataa gtataacgat attttagcaa 6660  
ataatatatt ttcaaacaaa acatgtttct ataatttttg tctaaccxaa aatttaggaa 6720  
tatgacctca attcttcaaa aagttagtaa aacaggttta aaaccccggt ataaatattt 6780  
ttgcctctga aacctatcaa attttcagat acaatcccgg tacacacaca tatcgcgaca 6840  
atcaatgtcg ctacgttctc gtcactgatt acaacgattt tgagctcgtt gagccagaat 6900  
tccgtcttcg ttggtatcag ggagatccga ctggtctcaa caatcagtat attctcaaga 6960  
tcattggacg acctgaatgc agcgagaaaa cattgagact tgaagtgaat ctttccacgt 7020  
gtgaaggtag attggaaact gcagagatga taggcgataa acgtcggaaa acaactttgt 7080  
tccagtggaa aaaatga 7097



## CE61773US.ST25

<210> 7  
<211> 3624  
<212> DNA  
<213> C. elegans

<400> 7  
atgtcaacaa tcaccagcca aaaagggata agattattaa ctgagagacg aggggataat 60  
tccctcatatc taactctcac tcttcactct ctctgctctt ctctcattt gtcttctttt 120  
tttgatattg gttgtggttt tttgtcaccg aataataaga atgctatgaa tacatctcac 180  
aattcatttt tctttttctt gcttctcttc cttttttcgt tctttttgcc gtttgccatt 240  
caactttttg gtaaattgcc aaattctaag aaaatgtggg ctttcccagc aattttgagc 300  
ataaatgtaa atctaatttc tagaaagttg atgggtcacag tgataccaaa aataataagt 360  
tctceatata ctcggacacg cctaccactg tacctctaca ctgtttccat cattatttcc 420  
tgctctttat tatactggaa tcttctttac tgcaaaaatt atgactgtgt cggtgagaag 480  
gaatttcgat ggggaagtac tcggcactta ctacagtact ttccggtgat agcagctccg 540  
attataatgg taatatcggt ttcttggtta ataattgcaa tatattattc aagtagttca 600  
tgtgttctta cattcaattt tatggaaatg ccatctgcag tactttgttc tctacttggt 660  
ggtattagtt ctgtaataga aattcatttt tccattgaag taaatcaagt tcaatggact 720  
gatcagtggg tactgtcatc tgtgggttta ccaatcaacg attgtttaaa aatcgatatt 780  
ttcaggggac ttcaatactt ttatgccttt tacatgctac aattgcgttc acacttcaat 840  
aatccttcca acatatttga atttccaatc ttcttcaaat cgatgaatca aaaatattat 900  
gtgaactgtg atatttactc ttgctcaatt catttcatga aaaagcaaaa gaaaatgagc 960  
ttttcacaag cacaagacgt atatcttcgt ctgaagcaag aaaaagaaga ggagaaacaa 1020  
cgagagcgag ccgaacgaga aaagcgaaat gagacgattg cagcgacaaa taaatcaaga 1080  
aagaagatga atcaggcatt ggcaaaaaga aataaaaaag gacaaccaa tctgaatgct 1140  
caaatggata tggcttccga tgaaaatatc ggtgccgacg gtgaacagaa gccttctcgg 1200  
ccgtttttga gaaaaggaca aggaacagca agatttagaa tggtagtttg tgcaaataca 1260  
aggcttatcg aaataatata tgaagttcag cctagaaaca acaaaacatc tgctgggtgca 1320

## CE61773US.ST25

cctccaacgt cggaactttc atctgcttca agtccttcta ttaatgttcc taggtttagt 1380  
ctgtcgaatg ctctcccgaa ctctgccga accgtggaca gtggaatatac aaatgaagac 1440  
gagaccggtc caccaaccac cgcattcttt ccaatggatc aaccatcatt gtcattcttcg 1500  
ccggaaaatc gtctaaatcc cgcaccttc gttgctgaag agcatggcca cagtggacag 1560  
cacgctgaag aagaagaaga caatgacacg gatgaagtat ctgcaatgcc ttcttttctg 1620  
cctgatgaac ctctgactct tgttaattca gatcatgaat tgtctgatga tgcttttaaag 1680  
tataaaaatg cagctgccga attcaaagct tttagagaga gaatggattc gatgagatca 1740  
gcttcaacaa tcacaacatc actggcaacg ccatcatctt gtgcaccatc aaactcctct 1800  
gagcctccta ctcggtctac accaattatg aacgatttag gcgttggccc aaataatcac 1860  
aattggccgt cttcaatgca agaattatca ggaatttctc tggaacacc acaggctcga 1920  
ccgcttgga gcaatagaat taatcagctt gttcgaagt aggctcaaac gggaataagc 1980  
cttttacaac accatgaaag acctactgtg accgccccat tgagacgaaa tgatatgatg 2040  
aactcatcac gacagaatcc acagaatgga aatgttcaag atgaaaatcg acccgagcac 2100  
gtttatgatc aaccaatata tgttcctgga tcatcactgg accgacagaa acttgaaatt 2160  
gaaattcgac gtcattgtaa cttgaacata caactgagag aactatttgc tcaattggat 2220  
tatgcagaag aatccgtgca caccacaaaa cgacagctcg aagaaaaaat ttccgaagtc 2280  
aataatttta agaaagaact gatagaagaa tttaagaaat gcaaaaaagg agttgaggaa 2340  
gaatttgaga agaagtgtga gaaaattaag gaagattatg atgaacttta cgagaaattg 2400  
aagagggatc aacgagatct tgaacgagat cagaagatat tgaagaaagg aacgggagaa 2460  
aggaataaag aattcacaga aacgatagcc actctccgag acaaatgaag agcatcagaa 2520  
accaagaatg cacaatatcg acaggatata cgtgttcgag acgaaaagct caagaaaaaa 2580  
gacgaggaaa tcgagaagct tcagaaagac ggaaaccggc taaagagcac tctacagact 2640  
ttagaaaagc gcgtaaaaca attacgtact gaaaaagaac gcgacgataa agaaaaggag 2700  
atgttcgaga aggttgcaat gaatcgaata acttcgaatc cagtgccacc agttttgaat 2760  
caaagtgttc caatttcgat aacatcaaat ggtccatcta gacatccatc atcatcttcg 2820  
ttgacaacat ttagaaaacc atctacatca aatcgagaaa gaggtgttag ttgggcagat 2880

CE61773US.ST25

gaaccaaattg aacaatcatt ggaagctgta ccacaggagt ttttgatgat gccagtcaaa 294  
 gaaatgccgg gaaaatttgg aaaatgcacg atctacagag attctcttgg agaaacatct 300  
 aaagtgcagg atacaatagc taacgggtctt cttttcgaat attccaatgg agatcttcga 306  
 tgggttaatc ggcagaacgc tgtaatatc tacatatccg cagttgataa aacagtcaga 312  
 attgatctcc ccacatacaa tatttcaatt attcatacat ttcaaaggca agttgaagta 318  
 cttcgtcctg gaaataacat aacattgata agtattaaac gacgagaagt tcgaactgat 324  
 ttgatttatc aaaacggaat gtataaaact gaaatcttca atagggacgg aagatatgtt 330  
 acgaaggatt ttagcaatca agaagtttcg agaaaataca atcccgttac acacacatat 336  
 cgcgacaatc aatgtcgcta cgttctcgtc actgattaca acgattttga gctcgttgag 342  
 ccagaattcc gtcttcggtg gtatcagga gatccgactg gtctcaaca tcagtatatatt 348  
 ctcaagatca ttggacgacc tgaatgcagc gagaaaacat tgagacttga agtgaatctt 354  
 tccacgtgtg aagggtacatt ggaaactgca gagatgatag gcgataaacg tcggaaaaca 360  
 actttgttcc agtggaaaaa atga 362

<210> 8  
 <211> 336  
 <212> PRT  
 <213> C. elegans

<400> 8

Met	Asn	Arg	Leu	Lys	Ser	Asp	Gln	Lys	Thr	Lys	Leu	Arg	Gln	Phe	Val
1				5					10					15	

Gln	Trp	Thr	Gln	Val	Thr	Glu	Ala	Val	Ser	Leu	Asn	Phe	Leu	Ala	Lys
			20					25					30		

Ala	Asn	Trp	Asn	Ile	Glu	Tyr	Ala	Met	Thr	Leu	Tyr	Phe	Asp	Asn	Pro
		35					40					45			

Asn	Leu	Phe	Ala	Gly	Ser	Thr	Pro	Gln	Pro	Ser	Val	Asp	Arg	Ser	Asn
	50					55					60				

Ile	Glu	Arg	Leu	Phe	Asn	Gln	Tyr	Val	Asp	Pro	Lys	Asp	Lys	Val	Gly
65					70					75					80

## CE61773US.ST25

Glu Lys Arg Met Gly Pro His Gly Ile Asn Arg Leu Leu Thr Asp Leu  
                     85                    90                    95

Gly Tyr Glu Ala Thr Asp Arg Arg Val Leu Val Leu Ala Trp Lys Phe  
                     100                    105                    110

Thr Ala Gln Thr Gln Cys Glu Phe Ser Leu Asp Glu Trp Val Lys Gly  
                     115                    120                    125

Met Thr Ala Leu Gln Ala Asp Thr Val Gln Asn Leu Arg Gln Arg Ile  
                     130                    135                    140

Asp Ser Ile Asn Ser Gly Leu Glu Ser Asp Lys Ala Lys Phe His Glu  
 145                    150                    155                    160

Leu Tyr Leu Phe Ala Phe Asn Tyr Ala Lys Ser Ala Ala Cys Arg Asn  
                     165                    170                    175

Leu Asp Leu Glu Thr Ala Ile Cys Cys Trp Asp Val Leu Phe Gly Gln  
                     180                    185                    190

Arg Ser Thr Ile Met Thr Gln Trp Ile Asp Phe Leu Trp Ala Gln Glu  
                     195                    200                    205

Asn Ala Ala Ala Ser Arg Leu Ala Gln Asn Val Gly Ala Ser Asn Ala  
                     210                    215                    220

Lys Gln Phe Lys Ser Val Trp Ile Ser Arg Asp Thr Trp Asn Leu Phe  
 225                    230                    235                    240

Trp Asp Phe Ile Leu Leu Ser Lys Pro Asp Leu Ser Asp Tyr Asp Asp  
                     245                    250                    255

Glu Gly Ala Trp Pro Val Leu Ile Asp Gln Phe Val Asp Tyr Cys Arg  
                     260                    265                    270

Glu Asn Leu Asn Tyr Pro Lys Pro Gly Asn Ala Ser Asn Asp Gln Gln  
                     275                    280                    285

CE61773US.ST25

Met Glu Thr Pro Lys Ile Ala Gln Lys Lys Pro Gly Ile Phe Tyr Phe  
 290 295 300

Asn Ser Asn Leu Gln Leu Ile Glu Phe Lys Leu Phe Gln Tyr Pro Met  
 305 310 315 320

Leu Lys Thr Ile Phe Lys Ile Thr Ile His Thr Ala Gly Thr Asn Arg  
 325 330 335

<210> 9

<211> 283

<212> PRT

<213> C. elegans

<400> 9

Met Asn Arg Leu Lys Ser Asp Gln Lys Thr Lys Ile Glu Arg Leu Phe  
 1 5 10 15

Asn Gln Tyr Val Asp Pro Lys Asp Lys Val Gly Glu Lys Arg Met Gly  
 20 25 30

Pro His Gly Ile Asn Arg Leu Leu Thr Asp Leu Gly Tyr Glu Ala Thr  
 35 40 45

Asp Arg Arg Val Leu Val Leu Ala Trp Lys Phe Thr Ala Gln Thr Gln  
 50 55 60

Cys Glu Phe Ser Leu Asp Glu Trp Val Lys Gly Met Thr Ala Leu Gln  
 65 70 75 80

Ala Asp Thr Val Gln Asn Leu Arg Gln Arg Ile Asp Ser Ile Asn Ser  
 85 90 95

Gly Leu Glu Ser Asp Lys Ala Lys Phe His Glu Leu Tyr Leu Phe Ala  
 100 105 110

Phe Asn Tyr Ala Lys Ser Ala Ala Cys Arg Asn Leu Asp Leu Glu Thr  
 115 120 125

## CE61773US.ST25

Ala Ile Cys Cys Trp Asp Val Leu Phe Gly Gln Arg Ser Thr Ile Met  
 130 135 140

Thr Gln Trp Ile Asp Phe Leu Trp Ala Gln Glu Asn Ala Ala Ala Ser  
 145 150 155 160

Arg Leu Ala Gln Asn Val Gly Ala Ser Asn Ala Lys Gln Phe Lys Ser  
 165 170 175

Val Trp Ile Ser Arg Asp Thr Trp Asn Leu Phe Trp Asp Phe Ile Leu  
 180 185 190

Leu Ser Lys Pro Asp Leu Ser Asp Tyr Asp Asp Glu Gly Ala Trp Pro  
 195 200 205

Val Leu Ile Asp Gln Phe Val Asp Tyr Cys Arg Glu Asn Leu Asn Tyr  
 210 215 220

Pro Lys Pro Gly Asn Ala Ser Asn Asp Gln Gln Met Glu Thr Pro Lys  
 225 230 235 240

Ile Ala Gln Lys Lys Pro Gly Ile Phe Tyr Phe Asn Ser Asn Leu Gln  
 245 250 255

Leu Ile Glu Phe Lys Leu Phe Gln Tyr Pro Met Leu Lys Thr Ile Phe  
 260 265 270

Lys Ile Thr Ile His Thr Ala Gly Thr Asn Arg  
 275 280

<210> 10  
 <211> 1010  
 <212> PRT  
 <213> C. elegans

<400> 10

Met Ser Met Glu Pro Arg Lys Lys Arg Asn Ser Ile Leu Lys Val Arg  
 1 5 10 15

## CE61773US.ST25

Gln Ala Val Glu Thr Ile Glu Glu Thr Val Met Asn Ser Gly Pro Ser  
 20 25 30

Ser Thr Thr Thr Asn Arg Arg Val Ser Phe His Asn Val Lys His Val  
 35 40 45

Lys Gln Tyr Asp Arg Asp His Gly Lys Ile Leu Asp Ala Thr Pro Val  
 50 55 60

Lys Glu Lys Ile Thr Asp Thr Ile Gly Ser Asp Gly Ile Leu Thr Pro  
 65 70 75 80

Arg Gly Gly Asn Met Asp Ile Ser Glu Ser Pro Ala Cys Thr Ser Ser  
 85 90 95

Phe Gln Val Phe Gly Gly Gly Asn Leu Asp Lys Thr Met Asp Met Ser  
 100 105 110

Leu Glu Thr Thr Ile Asn Glu Asn Asn Glu Thr Ala Arg Leu Phe Glu  
 115 120 125

Thr Thr Arg Asp Pro Thr Leu Leu Tyr Glu Lys Ile Val Glu Thr Thr  
 130 135 140

Thr Lys Val Thr Glu Arg Ile Val Ser Met Pro Leu Asp Asp Thr Leu  
 145 150 155 160

Ala Met Phe Asn Thr Thr Asn Gln Glu Asp Lys Asp Met Ser Val Asp  
 165 170 175

Arg Ser Val Leu Phe Thr Ile Pro Lys Val Pro Lys His Asn Ala Thr  
 180 185 190

Met Asn Arg Thr Ile Pro Met Asp Leu Asp Glu Ser Lys Ala Ala Gly  
 195 200 205

Gly Gln Cys Asp Glu Thr Met Asn Val Phe Asn Phe Thr Asn Leu Glu  
 210 215 220

## CE61773US.ST25

Ala Ala Glu Met Asp Thr Ser Lys Leu Asp Glu Asn Asn Thr Met Asn  
 225 230 235 240

Ala Ile Arg Ile Pro Ile Asn Ser Asn Val Met Pro Val Asp Met Asp  
 245 250 255

Ile Thr Glu His His Thr Leu Ile Glu Glu Lys Lys Asn Asp Thr Phe  
 260 265 270

Gly Pro Ser Gln Leu Met Asp Ile Ser Ala Pro Gln Val Gln Val Asn  
 275 280 285

Asp Thr Leu Ala Ile Phe Asn Ser Pro Arg Asp Ile Cys Asn Lys Gly  
 290 295 300

Leu Gly Val Pro Gln Asn Leu Ile Asn Ile Ala Ser Asn Val Val Pro  
 305 310 315 320

Val Asp Met Asp Ile Thr Asp Gln Ala Val Leu Asn Ala Glu Lys Lys  
 325 330 335

Asn Asp Gln Phe Glu Thr Ser Gln Leu Met Asp Ile Ser Ile Pro Lys  
 340 345 350

Val Leu Val Asn Asp Thr Met Ala Met Phe Asn Ser Pro Lys His Val  
 355 360 365

Ser Lys Ser Ser Met Asp Leu Glu Lys Thr Ile Glu Ala Ala Asp Lys  
 370 375 380

Ser Thr Lys Tyr Pro Ser Ile Ala Asp Glu Val Glu Asp Leu Asp Met  
 385 390 395 400

Asp Met Asp Ile Thr Glu Gln Gln Pro Cys Glu Ala Gly Asn Gln Gln  
 405 410 415

Asn Asp Gly Leu Gln Leu Gln Lys Glu Asp Leu Met Asp Ile Ser Val  
 420 425 430



## CE61773US.ST25

Ile Arg Asp Ser Pro Ala Val Asn Asp Thr Met Ala Val Phe Gln Ser  
 435 440 445

Pro Ala Arg Val Lys Ile Gly Ala Asn Asn Ser Ile Ile Asp Ser Gln  
 450 455 460

Lys Ser Ile Val Phe Gly Asp Glu Met Ser Ile Asp Glu Thr Gln Asn  
 465 470 475 480

Asp Gly Thr Leu Thr Leu Pro Lys Ser Asn Val Glu Val Thr Thr Thr  
 485 490 495

Asn Asp Val Tyr Thr Ser Leu Glu Arg Gln Glu Glu Asn Ala Ser Glu  
 500 505 510

Asn Val Ser Met Ile Asn Glu Ser Ser Val His Ser Glu Ile Asp Lys  
 515 520 525

Lys Ser Phe Met Leu Ile Glu Glu Glu Arg Ala Phe Met His Ser Ser  
 530 535 540

Met Ile Asp Val Ala Gln Lys Leu Glu Asp Asp Gly Ser Ser Lys Thr  
 545 550 555 560

Pro Val Ile Leu Ala Ser Gln Ser Ala Ser Leu Ala Thr Lys Glu Pro  
 565 570 575

Ser Ala Leu His Asn Ser Ser Ala Thr Leu Asn Asn Ser Met Glu Leu  
 580 585 590

Asp Asn Asn Thr Leu Leu Lys Thr Met Gln Ile Thr Thr Cys Glu Asp  
 595 600 605

Ile Ser Met Val His Glu Ser Ile Ala Val Glu Leu Asn Ser Asn Lys  
 610 615 620

Glu Gln Glu Gln Phe Gly Asp Glu Thr Leu Gln Lys Asn Asp Thr Ser  
 625 630 635 640

## CE61773US.ST25

Asn Thr Gly Ala Asn Phe Thr Phe Gln Gly His Asn Glu Thr Ser Gln  
 645 650 655

Ile Met Asn Asn Val Asp Ser Glu Ala Val Asn Thr Ser Lys Ile Ser  
 660 665 670

Thr Tyr Ser Ala Phe Asn Leu Ser Ile Asn Gln Ser Ile Ser Lys Arg  
 675 680 685

Arg Arg Ser Leu Leu Asn Ser Ala Arg Glu Ser Pro Arg Arg Val Ala  
 690 695 700

Leu Glu Asn Ser Ile Met Ser Met Asn Gly Gln Thr Met Glu Ala Leu  
 705 710 715 720

Thr Glu Tyr Arg Gln Asn Lys Thr Met Gln Thr Ser Gln Asp Ser Met  
 725 730 735

Pro Ser Met Ser Leu Asn Asp Ser Gly Arg Asp Ile Leu Ala Met Asn  
 740 745 750

Thr Ser Val Arg Ser Pro His Leu Asn Ser Ser Lys Thr Ala Ala Pro  
 755 760 765

Gly Thr Pro Ser Leu Met Ser Gln Asn Val Gln Leu Pro Pro Pro Ser  
 770 775 780

Pro Gln Phe Glu Met Pro Asp Phe Asp Pro Ala Val Val Asn Val Val  
 785 790 795 800

Tyr Leu Thr Ser Glu Asp Pro Ser Thr Glu Gln His Pro Glu Ala Leu  
 805 810 815

Lys Phe Gln Arg Ile Val Glu Asn Glu Lys Met Lys Val Gln His Glu  
 820 825 830

Ile Asp Ser Leu Asn Ser Thr Asn Gln Leu Ser Ala Glu Lys Ile Asp  
 835 840 845

## CE61773US.ST25

Met Leu Lys Thr Lys Glu Leu Leu Lys Phe Ser His Asp Glu Arg Glu  
 850 855 860  
 Ala Ile Met Ile Ala Arg Lys Asp Ala Glu Ile Lys Phe Leu Glu Leu  
 865 870 875 880  
 Arg Leu Lys Phe Ala Leu Glu Lys Lys Ile Glu Ser Asp Gln Glu Ile  
 885 890 895  
 Ala Glu Leu Glu Gln Gly Asn Ser Lys Met Ala Glu Gln Leu Arg Gly  
 900 905 910  
 Leu Asp Lys Met Ala Val Val Gln Lys Glu Leu Glu Lys Leu Arg Ser  
 915 920 925  
 Leu Pro Pro Ser Arg Glu Glu Ser Gly Lys Ile Arg Lys Glu Trp Met  
 930 935 940  
 Glu Met Lys Gln Trp Glu Phe Asp Gln Lys Met Lys Ala Leu Arg Asn  
 945 950 955 960  
 Val Arg Ser Asn Met Ile Ala Leu Arg Ser Glu Lys Asn Ala Leu Glu  
 965 970 975  
 Met Lys Val Ala Glu Glu His Glu Lys Phe Ala Gln Arg Asn Asp Leu  
 980 985 990  
 Lys Lys Ser Arg Met Leu Val Phe Ser Lys Ala Val Lys Lys Ile Val  
 995 1000 1005  
 Asn Phe  
 1010  
 <210> 11  
 <211> 1207  
 <212> PRT  
 <213> C. elegans  
 <400> 11

Met Ser Thr Ile Thr Ser Gln Lys Gly Ile Arg Leu Leu Thr Glu Arg

CE61773US.ST25

1	5	10	15
Arg Gly Asp Asn Ser Leu Ile Leu Thr Leu Thr Leu His Ser Leu Cys	20	25	30
Ser Ser Pro His Leu Ser Ser Phe Phe Asp Ile Gly Cys Gly Phe Leu	35	40	45
Ser Pro Asn Asn Lys Asn Ala Met Asn Thr Ser His Asn Ser Phe Phe	50	55	60
Phe Phe Leu Leu Leu Phe Leu Phe Ser Phe Phe Leu Pro Phe Ala Ile	65	70	75
Gln Leu Phe Gly Lys Leu Pro Asn Ser Lys Lys Met Trp Ala Phe Pro	85	90	95
Ala Ile Leu Ser Ile Asn Val Asn Leu Ile Ser Arg Lys Leu Met Val	100	105	110
Thr Val Ile Pro Lys Ile Ile Ser Ser Pro Tyr Pro Arg Thr Arg Leu	115	120	125
Pro Leu Tyr Leu Tyr Thr Val Ser Ile Ile Ile Ser Cys Ser Leu Leu	130	135	140
Tyr Trp Asn Leu Leu Tyr Cys Lys Asn Tyr Asp Cys Val Val Glu Lys	145	150	155
Glu Phe Arg Trp Gly Ser Thr Arg His Leu Leu Gln Tyr Phe Pro Val	165	170	175
Ile Ala Ala Pro Ile Ile Met Val Ile Ser Phe Ser Trp Leu Ile Ile	180	185	190
Ala Ile Tyr Tyr Ser Ser Ser Ser Cys Val Leu Thr Phe Asn Phe Met	195	200	205
Glu Met Pro Ser Ala Val Leu Cys Ser Leu Leu Gly Gly Ile Ser Ser			

210                      215                      220

Cys Ala Asn Thr Arg Leu Ile Glu Ile Ile Tyr Glu Val Gln Pro Arg

## CE61773US.ST25

420	425	430
Asn Asn Lys Thr Ser Ala Gly Ala Pro Pro Thr Ser Glu Leu Ser Ser	435	440
Ala Ser Ser Pro Ser Ile Asn Val Pro Arg Phe Ser Leu Ser Asn Ala	450	460
Leu Pro Asn Ser Ala Arg Thr Val Asp Ser Gly Ile Ser Asn Glu Asp	465	470
Glu Thr Arg Pro Pro Thr Thr Ala Ser Leu Pro Met Asp Gln Pro Ser	485	490
Leu Ser Ser Ser Pro Glu Asn Arg Leu Asn Pro Ala Pro Ser Val Ala	500	510
Glu Glu His Gly His Ser Gly Gln His Ala Glu Glu Glu Glu Asp Asn	515	520
Asp Thr Asp Glu Val Ser Ala Met Pro Ser Phe Val Pro Asp Glu Pro	530	535
Ser Thr Leu Val Asn Ser Asp His Glu Leu Ser Asp Asp Ala Leu Lys	545	550
Tyr Lys Asn Ala Ala Ala Glu Phe Lys Ala Phe Glu Arg Arg Met Asp	565	570
Ser Met Arg Ser Ala Ser Thr Ile Thr Thr Ser Leu Ala Thr Pro Ser	580	585
Ser Cys Ala Pro Ser Asn Ser Ser Glu Pro Pro Thr Arg Ser Thr Pro	595	600
Ile Met Asn Asp Leu Gly Val Gly Pro Asn Asn His Asn Trp Pro Ser	610	615
Ser Met Gln Glu Leu Ser Gly Ile Ser Leu Glu Thr Pro Gln Ala Arg		

CE61773US.ST25

625		630		635		640									
Pro	Leu	Gly	Ser	Asn	Arg	Ile	Asn	Gln	Leu	Val	Arg	Ser	Glu	Ala	Gln
				645					650					655	
Thr	Gly	Ile	Ser	Leu	Leu	Gln	His	His	Glu	Arg	Pro	Thr	Val	Thr	Ala
			660					665					670		
Pro	Leu	Arg	Arg	Asn	Asp	Met	Met	Asn	Ser	Ser	Arg	Gln	Asn	Pro	Gln
		675					680					685			
Asn	Gly	Asn	Val	Gln	Asp	Glu	Asn	Arg	Pro	Glu	His	Val	Tyr	Asp	Gln
	690					695					700				
Pro	Ile	His	Val	Pro	Gly	Ser	Ser	Leu	Asp	Arg	Gln	Lys	Leu	Glu	Ile
705					710					715					720
Glu	Ile	Arg	Arg	His	Arg	Asn	Leu	Asn	Ile	Gln	Leu	Arg	Asp	Thr	Ile
				725					730					735	
Ala	His	Leu	Asp	Tyr	Ala	Glu	Glu	Ser	Val	His	Thr	Thr	Lys	Arg	Gln
			740					745					750		
Leu	Glu	Glu	Lys	Ile	Ser	Glu	Val	Asn	Asn	Phe	Lys	Lys	Glu	Leu	Ile
		755					760					765			
Glu	Glu	Phe	Lys	Lys	Cys	Lys	Lys	Gly	Val	Glu	Glu	Glu	Phe	Glu	Lys
	770					775					780				
Lys	Phe	Glu	Lys	Ile	Lys	Glu	Asp	Tyr	Asp	Glu	Leu	Tyr	Glu	Lys	Leu
785					790					795					800
Lys	Arg	Asp	Gln	Arg	Asp	Leu	Glu	Arg	Asp	Gln	Lys	Ile	Leu	Lys	Lys
				805					810					815	
Gly	Thr	Gly	Glu	Arg	Asn	Lys	Glu	Phe	Thr	Glu	Thr	Ile	Ala	Thr	Leu
			820					825					830		
Arg	Asp	Lys	Leu	Arg	Ala	Ser	Glu	Thr	Lys	Asn	Ala	Gln	Tyr	Arg	Gln

CE61773US.ST25

835

840

845

Asp Ile Arg Val Arg Asp Glu Lys Leu Lys Lys Lys Asp Glu Glu Ile  
 850 855 860

Glu Lys Leu Gln Lys Asp Gly Asn Arg Leu Lys Ser Thr Leu Gln Thr  
 865 870 875 880

Leu Glu Lys Arg Val Lys Gln Leu Arg Thr Glu Lys Glu Arg Asp Asp  
 885 890 895

Lys Glu Lys Glu Met Phe Ala Lys Val Ala Met Asn Arg Lys Thr Ser  
 900 905 910

Asn Pro Val Pro Pro Val Leu Asn Gln Ser Val Pro Ile Ser Ile Thr  
 915 920 925

Ser Asn Gly Pro Ser Arg His Pro Ser Ser Ser Ser Leu Thr Thr Phe  
 930 935 940

Arg Lys Pro Ser Thr Ser Asn Arg Glu Arg Gly Val Ser Trp Ala Asp  
 945 950 955 960

Glu Pro Asn Glu Gln Ser Leu Glu Ala Val Pro Gln Glu Phe Leu Met  
 965 970 975

Met Pro Val Lys Glu Met Pro Gly Lys Phe Gly Lys Cys Thr Ile Tyr  
 980 985 990

Arg Asp Ser Leu Gly Glu Thr Ser Lys Val Thr Asp Thr Ile Ala Asn  
 995 1000 1005

Gly Leu Leu Phe Glu Tyr Ser Asn Gly Asp Leu Arg Trp Val Asn  
 1010 1015 1020

Arg Gln Asn Ala Val Asn Ile Tyr Ile Ser Ala Val Asp Lys Thr  
 1025 1030 1035

Val Arg Ile Asp Leu Pro Thr Tyr Asn Ile Ser Ile Ile His Thr



```
<210> 12
<211> 780
<212> DNA
<213> homo sapiens
```

## CE61773US.ST25

```

<400> 12
atgaacaagt tgaaatcatc gcagaaggat aaagttcgtc agtttatgat cttcacacaa 60
tctagtgaaa aaacagcagt aagttgtctt tctcaaatg actggaagtt agatgttgca 120
acagataatt ttttccaaaa tcctgaactt tatatacgag agagtgtaaa aggatcattg 180
gacaggaaga agttagaaca gctgtacaat agatacaaag accctcaaga tgagaataaa 240
attggaatag atggcataca gcagttctgt gatgacctgg cactcgatcc agccagcatt 300
agtgtgttga ttattgctg gaagttcaga gcagcaacac agtgcgagtt ctccaaacag 360
gagttcatgg atggcatgac agaattagga tgtgacagca tagaacaact aaaggcccag 420
ataccaaga tggaacaaga attgaaagaa ccaggacgat ttaaggattt ttaccagttt 480
acttttaatt ttgcaaagaa tccaggacaa aaaggattag atctagaaat ggccattgcc 540
tactggaact tagtgcttaa tggaagattt aaattcttag acttatggaa taaatttttg 600
ttggaacatc ataaacgatc aatacaaaaa gacacttgga atcttctttt agacttcagt 660
acgatgattg cagatgacat gtctaattat gatgaagaag gagcatggcc tgttcttatt 720
gatgactttg tggaatttgc acgccctcaa attgctggga caaaaagtac aacagtgtag 780

```

```

<210> 13
<211> 259
<212> PRT
<213> homo sapiens

```

```

<400> 13

```

```

Met Asn Lys Leu Lys Ser Ser Gln Lys Asp Lys Val Arg Gln Phe Met
1           5           10           15

```

```

Ile Phe Thr Gln Ser Ser Glu Lys Thr Ala Val Ser Cys Leu Ser Gln
20           25           30

```

```

Asn Asp Trp Lys Leu Asp Val Ala Thr Asp Asn Phe Phe Gln Asn Pro
35           40           45

```

```

Glu Leu Tyr Ile Arg Glu Ser Val Lys Gly Ser Leu Asp Arg Lys Lys
50           55           60

```

```

Leu Glu Gln Leu Tyr Asn Arg Tyr Lys Asp Pro Gln Asp Glu Asn Lys

```

## CE61773US.ST25

65                                      70                                      75                                      80  
 Ile Gly Ile Asp Gly Ile Gln Gln Phe Cys Asp Asp Leu Ala Leu Asp  
    85                                      90                                      95  
 Pro Ala Ser Ile Ser Val Leu Ile Ile Ala Trp Lys Phe Arg Ala Ala  
    100                                      105                                      110  
 Thr Gln Cys Glu Phe Ser Lys Gln Glu Phe Met Asp Gly Met Thr Glu  
    115                                      120                                      125  
 Leu Gly Cys Asp Ser Ile Glu Gln Leu Lys Ala Gln Ile Pro Lys Met  
    130                                      135                                      140  
 Glu Gln Glu Leu Lys Glu Pro Gly Arg Phe Lys Asp Phe Tyr Gln Phe  
 145                                      150                                      155                                      160  
 Thr Phe Asn Phe Ala Lys Asn Pro Gly Gln Lys Gly Leu Asp Leu Glu  
    165                                      170                                      175  
 Met Ala Ile Ala Tyr Trp Asn Leu Val Leu Asn Gly Arg Phe Lys Phe  
    180                                      185                                      190  
 Leu Asp Leu Trp Asn Lys Phe Leu Leu Glu His His Lys Arg Ser Ile  
    195                                      200                                      205  
 Pro Lys Asp Thr Trp Asn Leu Leu Leu Asp Phe Ser Thr Met Ile Ala  
    210                                      215                                      220  
 Asp Asp Met Ser Asn Tyr Asp Glu Glu Gly Ala Trp Pro Val Leu Ile  
 225                                      230                                      235                                      240  
 Asp Asp Phe Val Glu Phe Ala Arg Pro Gln Ile Ala Gly Thr Lys Ser  
    245                                      250                                      255  
 Thr Thr Val

&lt;210&gt; 14

CE61773US.ST25

&lt;211&gt; 258

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

Ser Lys Gln Glu Phe Met Asp Gly Met Thr Glu Leu Gly Cys Asp Ser  
 1 5 10 15

Ile Glu Gln Leu Lys Ala Gln Ile Pro Lys Met Glu Gln Glu Leu Lys  
 20 25 30

Glu Pro Gly Arg Phe Lys Asp Phe Tyr Gln Phe Thr Phe Asn Phe Ala  
 35 40 45

Lys Asn Pro Gly Gln Lys Gly Leu Asp Leu Glu Asp Arg Lys Lys Leu  
 50 55 60

Glu Gln Leu Tyr Asn Arg Tyr Lys Asp Pro Gln Asp Glu Asn Lys Ile  
 65 70 75 80

Gly Ile Asp Gly Ile Gln Gln Phe Cys Asp Asp Leu Ala Leu Asp Pro  
 85 90 95

Ala Ser Ile Ser Val Leu Ile Ile Ala Trp Lys Phe Arg Ala Ala Thr  
 100 105 110

Gln Cys Glu Phe Ser Lys Gln Glu Phe Met Asp Gly Met Thr Glu Leu  
 115 120 125

Gly Cys Asp Ser Ile Glu Gln Leu Lys Ala Gln Ile Pro Lys Met Glu  
 130 135 140

Gln Glu Leu Lys Glu Pro Gly Arg Phe Lys Asp Phe Tyr Gln Phe Thr  
 145 150 155 160

Phe Asn Phe Ala Lys Asn Pro Gly Gln Lys Gly Leu Asp Leu Glu Met  
 165 170 175

Ala Ile Ala Tyr Trp Asn Leu Val Leu Asn Gly Arg Phe Lys Phe Leu  
 180 185 190

CE61773US.ST25

Asp Leu Trp Asn Lys Phe Leu Leu Glu His His Lys Arg Ser Ile Pro  
 195 200 205

Lys Asp Thr Trp Asn Leu Leu Leu Asp Phe Ser Thr Met Ile Ala Asp  
 210 215 220

Asp Met Ser Asn Tyr Asp Glu Glu Gly Ala Trp Pro Val Leu Ile Asp  
 225 230 235 240

Asp Phe Val Glu Phe Ala Arg Pro Gln Ile Ala Gly Thr Lys Ser Thr  
 245 250 255

Thr Val

<210> 15  
 <211> 19  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> T7 polymerase promoter sequence (example 1)

<400> 15  
 taatacgact cactatagg 19

<210> 16  
 <211> 19  
 <212> DNA  
 <213> artificial sequence

<220>  
 <223> T3 polymerase promoter sequence

<400> 16  
 aattaaccct cactaaagg 19

<210> 17  
 <211> 19  
 <212> DNA  
 <213> artificial sequence

<220>

## CE61773US.ST25

<223> oligonucleotide for PCR amplification (example 3)

<400> 17  
tcaatcagta tgtcgaccc 19

<210> 18  
<211> 19  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 3)

<400> 18  
ggaagaaatt ggggaaaca 19

<210> 19  
<211> 19  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 3)

<400> 19  
atcgagcgcc tcttcaatc 19

<210> 20  
<211> 19  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 3)

<400> 20  
tggtgtctcc atttgctga 19

<210> 21  
<211> 19  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 4)

<400> 21  
atctgaagat ccgtccact 19

## CE61773US.ST25

<210> 22  
<211> 19  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 4)

<400> 22  
atgcacaatg ggtatTTTTT 19

<210> 23  
<211> 23  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 5; forward primer to generate dsRNA 305A12)

<400> 23  
ttcgtctcga acacgtatat cct 23

<210> 24  
<211> 23  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 5; reverse primer to generate dsRNA 305A12)

<400> 24  
gaaagaagat gaatcaggca ttg 23

<210> 25  
<211> 23  
<212> DNA  
<213> artificial sequence

<220>  
<223> oligonucleotide for PCR amplification (example 5; forward primer to generate dsRNA 341G5)

<400> 25  
ctgcaaaaat tatgactgtg tcg 23

CE61773US.ST25

&lt;210&gt; 26

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; artificial sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide for PCR amplification (example 5; reverse primer to generate dsRNA 341G5)

&lt;400&gt; 26

agcattcaga tttggttgtc c

21



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**